


08-18-00

A

08/17/00



1c883 U.S. PTO

UTILITY PATENT APPLICATION TRANSMITTAL

Submit an original and a duplicate for fee processing
(Only for new nonprovisional applications under 37 CFR §1.53(b))

ADDRESS TO:

Commissioner of Patents and Trademarks
Box Patent Application
Washington, D.C. 20231

Attorney Docket No. 205970

First Named Inventor Arnd BAUMANN

Express Mail No. EL643535335US

1c857 U.S. PTO

09/640582



08/17/00

APPLICATION ELEMENTS

1. ☒ Utility Transmittal Form
2. ☒ Specification (including claims and abstract) [Total Pages 47]
3. ☒ Drawings [Total Sheets 18]
4. ☐ Combined Declaration and Power of Attorney [Total Pages]
 - a. ☐ Newly executed
 - b. ☐ Copy from prior application
 - i. ☐ Deletion of Inventor(s) Signed statement attached deleting inventor(s) named in the prior application
5. ☐ Incorporation by Reference: The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
6. ☐ Microfiche Computer Program
7. ☒ Nucleotide and/or Amino Acid Sequence Submission
 - a. ☐ Computer Readable Copy
 - b. ☒ Paper Copy
 - c. ☐ Statement verifying above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet and document(s))
9. ☐ Power of Attorney
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure Statement (IDS)
 - ☐ Form PTO-1449
 - ☐ Copies of References
12. ☐ Preliminary Amendment
13. ☒ Return Receipt Postcard (Should be specifically itemized)
14. ☐ Small Entity Statement(s)
 - ☐ Enclosed
 - ☐ Statement filed in prior application; status still proper and desired
15. ☐ Certified Copy of Priority Document(s)
16. ☒ Other: Patent Application Cover Sheet

17. If a **CONTINUING APPLICATION**, check appropriate box and supply the requisite information in (a) and (b) below:
- (a) ☐ Continuation ☐ Divisional ☒ Continuation-in-part of prior application Serial No. PCT/EP99/00942 filed February 12, 1999.
Prior application information: Examiner ; Group Art Unit:
- (b) Preliminary Amendment: Relate Back - 35 USC §120. The Commissioner is requested to amend the specification by inserting the following sentence before the first line:
"This is a ☒ continuation-in-part ☐ divisional of copending application(s)
☐ Application No. , filed on
☒ International Application No. PCT/EP99/00942, filed on February 12, 1999, and which designates the U.S."

APPLICATION FEES

BASIC FEE				\$690.00
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total Claims	46 -20=	26	x \$18.00	\$468.00
Independent Claims	5 -3=	2	x \$78.00	\$156.00
<input type="checkbox"/> Multiple Dependent Claims(s) if applicable			+\$260.00	\$
Total of above calculations =				\$1,314.00
Reduction by 50% for filing by small entity =				\$(0.00)
<input type="checkbox"/> Assignment fee if applicable			+\$40.00	\$
TOTAL =				\$1,314.00

UTILITY PATENT APPLICATION TRANSMITTAL

Attorney Docket No. 205970

19. ☒ Please charge my Deposit Account No. 12-1216 in the amount of \$1,314.00.
20. ☐ A check in the amount of \$ is enclosed.
21. The Commissioner is hereby authorized to credit overpayments or charge any additional fees of the following types to Deposit Account No. 12-1216:
- a. ☒ Fees required under 37 CFR §1.16.
- b. ☒ Fees required under 37 CFR §1.17.
22. ☒ The Commissioner is hereby generally authorized under 37 CFR §1.136(a)(3) to treat any future reply in this or any related application filed pursuant to 37 CFR §1.53 requiring an extension of time as incorporating a request therefor, and the Commissioner is hereby specifically authorized to charge Deposit Account No. 12-1216 for any fee that may be due in connection with such a request for an extension of time.

23. CORRESPONDENCE ADDRESS

☒ Customer Number: 23460**23460**

PATENT TRADEMARK OFFICE



, Reg. No.

Leydig, Voit & Mayer, Ltd.
Two Prudential Plaza, Suite 4900
180 North Stetson
Chicago, Illinois 60601-6780
(312) 616-5600 (telephone)
(312) 616-5700 (facsimile)

Name	Carol Larcher, Registration No. 35,243
Signature	
Date	August 17, 2000

Certificate of Mailing Under 37 CFR §1.10

I hereby certify that this Utility Patent Application Transmittal and all accompanying documents are being deposited with the United States Postal Service "Express Mail Post Office To Addressee" Service under 37 CFR §1.10 on the date indicated below and is addressed to: Commissioner of Patents and Trademarks, Box Patent Application, Washington, D.C. 20231.

<i>Iring Mikiticuk</i>	<i>J. Mikiticuk</i>	August 17, 2000
Name of Person Signing	Signature	Date

PATENT APPLICATION

Invention Title:

SEQUENCES OF AN I_H ION CHANNEL AND USE THEREOF

Inventors:

Arnd BAUMANN	Germany	Juelich	Germany
INVENTOR'S NAME	CITIZENSHIP	CITY OF RESIDENCE	STATE or FOREIGN COUNTRY

Wolfgang BONIGK	Germany	Juelich	Germany
INVENTOR'S NAME	CITIZENSHIP	CITY OF RESIDENCE	STATE or FOREIGN COUNTRY

Renate GAUSS	Germany	Juelich	Germany
INVENTOR'S NAME	CITIZENSHIP	CITY OF RESIDENCE	STATE or FOREIGN COUNTRY

Alexander SCHOLTEN	Germany	Dormagen	Germany
INVENTOR'S NAME	CITIZENSHIP	CITY OF RESIDENCE	STATE or FOREIGN COUNTRY

Reinhard SEIFERT	Germany	Aachen	Germany
INVENTOR'S NAME	CITIZENSHIP	CITY OF RESIDENCE	STATE or FOREIGN COUNTRY

Benjamin KAUPP	Germany	Aachen	Germany
INVENTOR'S NAME	CITIZENSHIP	CITY OF RESIDENCE	STATE or FOREIGN COUNTRY

Be it known that the inventors listed above have invented a certain new and useful invention with the title shown above of which the following is a specification.

Sequences of an I_h ion channel and use thereof

The present invention relates to a nucleic acid, preferably a DNA, comprising at least part of the sequence of an I_h ion channel. Said sequence may e.g. be derived from a human DNA, a rat DNA, a bovine DNA, a *Drosophila melanogaster* DNA or a sea urchin DNA. Furthermore, the present invention relates to an mRNA molecule which contains the corresponding sequences. The invention further relates to a polypeptide or protein comprising the corresponding derived amino acid sequence.

Furthermore, the invention relates to the use of one or more of the above-mentioned sequences in a screening and/or diagnosing method and to the kits required therefor.

Lastly, the invention relates to the use of one or more of the above-mentioned sequences for the treatment and/or prophylaxis of cardiovascular disorders and sleep disturbances.

The many different functions of the nerve system are substantially determined by finely adjusted interactions between the intrinsic characteristics of the neurons and the synaptic connections. The electrophysiological characteristics inherent to the neurons and synapses are, in turn, determined by the localization and density of the voltage- and ligand-controlled ion channels which regulate the flow of ion currents across the neuronal plasma membrane and which are controlled by a great number of transmitter substances and intracellular messenger systems (Hille, 1992).

With regard to the specific activity expected of the neuronal elements, it is not astonishing that neurons have a great repertory of ion channels, including the classic

channels that produce voltage-dependent sodium (Na^+) and potassium (K^+) currents during an action potential (Hodgkin and Huxley, 1952) and also a number of unusual ion conductances (Linás, 1988).

An unusual intrinsic mechanism which had originally been discovered by Ito and colleagues (Araki et al., 1962; Ito and Oshima, 1965) in motoneurons of cats turned out to be a slow relaxation of the potential change induced by hyperpolarizing current, resulting in a non-ohmic behavior of the current/voltage (I/V) relationship in hyperpolarizing direction. The underlying time-dependent membrane current was first characterized in photoreceptors of the rods as cesium (Cs^+)-sensitive inward current which is triggered by hyperpolarization and may depolarize the membrane. This leads to the typical sequence of an initial transient hyperpolarization by exposure, followed by a slow depolarization (Attwell and Wilson, 1980; Bader et al., 1982; Bader et al., 1979; Fain et al., 1978).

The current in the photoreceptors was designated as I_h because it is activated by hyperpolarization. At about the same time a similar ion current was discovered in the heart, in the pacemaker cells of the sinus node and in the Purkinje fibers of the mammalian heart (Brown and DiFrancesco, 1980; Brown et al., 1979; DiFrancesco, 1981a; DiFrancesco, 1981b; Yanagihara and Irisawa, 1980), and it became clear that the slow inward current is accompanied by sodium and potassium ions. This current was called "funny" current (I_f) to emphasize its unusual behavior, i.e., the fact that an inward current is concerned which is activated by hyperpolarization and, oddly enough, was similar to the previously described K^+ conductance I_{K2} . There is a growing interest in said current because it participates, for instance, in the generation and control of spontaneous activity of the heart.

004700" 28504960

Further evidence of the presence of a corresponding current in central neurons was found, and it was mentioned by Halliwell and Adams (1982) for the first time. They observed a slow inward current, which was designated as "queer" current (I_q), in pyramidal cells of the hippocampus after hyperpolarization. Subsequently, currents with similar characteristics were found in a great number of neuronal and non-neuronal cells, and said hyperpolarization-activated current was finally recognized as an omnipresent phenomenon in cells of the nerve system. The designation as " I_h " is now accepted as a term for describing said current.

Although it was first assumed that the activity of the respective I_h channels is not modulated, more and more data show that the I_h channels are important targets for neurotransmitters and messenger systems, which emphasizes their important physiological role in the control of cellular electrical activities.

In the meantime it has become known that I_h significantly contributes to the rest potential, limits an excessive hyperpolarization, determines the form of action patterns (firing patterns) and takes part in the generation of rhythmic oscillations of the membrane potential. I_h currents have a few special characteristics that distinguish the same from other voltage-controlled ion channels. Like voltage-controlled Na^+ , Ca^{2+} and specific K^+ currents, they have a steep voltage-dependence curve and activate with a sigmoidal time course; they are however activated by hyperpolarization and deactivate by sigmoidal kinetics.

The activation in negative potentials and the blockage by Cs^+ ions reminds of inwardly rectifying K^+ channels. However, many characteristics of I_h clearly differ from that K^+ channel family: The activation kinetics is slower, the activation range is more positive and is independent of the extracellular K^+ concentration, conductance is substantially resistant to extracellular Ba^{2+} ions and the I_h channels are permeable not only to K^+ ions,

but also to Na^+ ions. In contrast to other cation channels, such as ligand-controlled cation channels, the I_h channels are very selective for Na^+ and K^+ ions and have a steep voltage-dependent control.

Of particular importance to the present research work is the participation of the I_h channels in the pacemaker function in the cardiac muscle. The pacemaker activity in the heart is due to specialized myocytes that are located in specific regions of the heart (*sinus venosus*) and are characterized by their ability to beat spontaneously even if separated from the rest of the cardiac muscle. In pacemaker cells of the sinus node in mammals, the spontaneous activity follows from a typical phase of their action potential, the slow diastolic depolarization. During said phase, which corresponds to the diastole of the cardiac contraction cycle, the membrane depolarizes again at a slow pace after termination of the action potential until the threshold value for the generation of a new action potential is reached. Thus the diastolic depolarization is responsible for the initiation of the rhythmic behavior and characterizes action potentials of the sinus node and other spontaneously active cardiocytes.

Apart from the generation of a rhythmic activity, the diastolic (or pacemaker) depolarization takes part in the control of the heartbeat frequency by autonomous neurotransmitters. It is known that the stimulation of the sympathetic and parasympathic nerve system leads to an acceleration and deceleration of the heartbeat.

It has become known in the meantime that the I_h channels take part in this pacemaker function. The I_h current of the sinus node is an unspecific cation current, normally accompanied by Na^+ and K^+ , which after hyperpolarization slowly activates in a voltage range encompassing that of the diastolic depolarization. The I_h features are well suited for producing a depolarization process as a reaction to a hyperpolarization in a voltage range in which the I_h channel is activated.

So far, however, it has not been possible to identify sequences of genes coding for I_h ion channels. Furthermore, channel protein has so far not been available in a sufficient amount for characterizing the same biochemically. Finally, the pharmacological characterization of I_h channels has so far been extremely difficult because the I_h currents were identified on whole cells, which additionally exhibit K^+ - and Na^+ -selective conductivities, and were experimentally isolated from the other currents.

It has therefore been the object of the present invention to indicate the nucleic acid, to show its possible applications, and to provide the protein in a functional state and in a sufficient amount for biochemical analyses and pharmaceutical applications.

Said object is achieved by the subject matter of the independent claims. Advantageous developments are indicated in the dependent claims.

The terms used hereinafter shall have the following meanings:

" I_h ion channel" is here to stand for those ion channels that (1) open by hyperpolarization and are closed at more positive voltage values ($V_m \geq -10$ mV); (2) whose activation and deactivation take place with a relatively slow sigmoidal time course; (3) conduct not only K^+ ions, but also Na^+ ions; (4) are almost entirely blocked by 0.1 – 3 mM extracellular Cs^+ and (5) are directly modulated by cyclic nucleotides, in particular cyclo AMP and cyclo GMP.

"Stringent conditions" means hybridization with 0.1-5 x SSC, preferably 1-2 x SSC, at 60-70°C, preferably 65°C.

"Conditions of low stringency" means hybridization at 0.1-5 x SSC, preferably 1-2 x SSC at 50-60°C, preferably at 55°C.

"Parts" of the I_h ion channel means a section of the protein sequence suited as antigenic determinant, for example, a section of at least 6 amino acids. Sections that occur in the form of domains, such as the sections S1, S2, etc. as indicated in Fig. 1A, are also regarded as parts. This encompasses sections of the ion channel that derive from the DNA sequences indicated in SEQ ID NO 1 to 15 using the IUPAC code, namely by way of amino acid exchanges, deletions and additions, while maintaining the biological function.

"Part" thereof in connection with the nucleic acid means a fragment having a length of at least 6 nucleotides, preferably 12 nucleotides, particularly preferably a length of 18 nucleotides. The part is suited for hybridizing via oligonucleotide hybridization specifically (selectively) with the corresponding total sequence. Thus a "part" of the nucleic acid is a section from the sequences according to SEQ ID NO 1 to 15 that is suited for selectively hybridizing with one of the said sequences.

"Selectively" (specifically) means that under suitable hybridization conditions a nucleic acid only hybridizes with one nucleic acid as is indicated by one of the sequences according to SEQ ID NO 1 to 15, whereas it does not hybridize with another nucleic acid of the respective host organism with which it is normally associated.

"Homology" as is here used is calculated as follows: The amino acids are counted in the sequences or sequence sections to be compared that are either identical or similar at the respective position. This number is divided by the total number of the amino acid residues and multiplied by 100. This yields a percentage of the sequence similarity or homology. This is illustrated by the sample given below:

TWALFKALSHMLCIGYGKFPPQS
PDAFWWAVVTMTTVGYGDMTPVG

The total number of the positions to be compared with one another is 23 residues; there are 7 identically and 6 similarly occupied amino acid positions. That is why the homology $(7 + 6)/23 \times 100 = 56.5\%$. An exchange of similar amino acids is also designated as a conservative exchange (cf. Dayhoff et al., 1978).

According to claim 1 there is provided a nucleic acid which comprises at least a part of the sequence of an I_h ion channel. The nucleic acid complementary thereto is also regarded as an inventive embodiment. Said nucleic acid may preferably be derived from a human DNA and is then in particular characterized by the sequences according to SEQ ID NO 1, SEQ ID NO 10, SEQ ID NO 11 and SEQ ID NO 15.

Advantageously, the sequence may also be derived from a rat DNA and is then in particular characterized by the SEQ ID NO 2 and SEQ ID NO 8, SEQ ID NO 9, SEQ ID NO 13 and SEQ ID NO 14.

In a further preferred embodiment, the sequence may be derived from a bovine DNA and is then characterized by the sequences according to SEQ ID NO 3 and SEQ ID NO 6, SEQ ID NO 7 and SEQ ID NO 12.

Furthermore, the sequence may preferably be derived from a sea urchin DNA, and it is then preferably characterized by the sequence SEQ ID NO 4.

Furthermore, the DNA may preferably be derived from *Drosophila melanogaster*. The complete sequence is then in accordance with SEQ ID NO 5.

Insert 1

0040502.0040502

Insert 1:

The above isolated or purified nucleic acid molecules also can be characterized in terms of "percentage of sequence identity." In this regard, a given nucleic acid molecule as described above can be compared to a nucleic acid molecule encoding a corresponding gene (i.e., the reference sequence) by optimally aligning the nucleic acid sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence, which does not comprise additions or deletions, for optimal alignment of the two sequences. The percentage of sequence identity is calculated by determining the number of positions at which the identical nucleic acid base occurs in both sequences, i.e., the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity. Optimal alignment of sequences for comparison may be conducted by computerized implementations of known algorithms (e.g., GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI, or BlastN and BlastX available from the National Center for Biotechnology Information, Bethesda, MD), or by inspection. Sequences are typically compared using BESTFIT or BlastN with default parameters.

"Substantial sequence identity" means that at least 75%, preferably at least 80%, more preferably at least 90%, and most preferably at least 95% of the sequence of a given nucleic acid molecule is identical to a given reference sequence. Typically, two polypeptides are considered to be substantially similar if at least 40%, preferably at least 60%, more preferably at least 90%, and most preferably at least 95% of the amino acids of which the polypeptides are comprised are identical to or represent conservative substitutions of the amino acids of a given reference sequence.

One of ordinary skill in the art will appreciate, however, that two polynucleotide sequences can be substantially different at the nucleic acid level, yet encode substantially similar, if not identical, amino acid sequences, due to the

0040527 084700

degeneracy of the genetic code. The present invention is intended to encompass such polynucleotide sequences.

007700" 22504350

A particularly preferred embodiment comprises sequences that exhibit a homology of at least 80% to one of the sequences with the SEQ ID NO 1 to 15. In a further preferred embodiment the sequence exhibits a homology of at least 90% to one of the sequences designated by SEQ ID NO 1 to 15.

It hybridizes in a particularly preferred manner under low stringent conditions and even more preferably under conditions of high stringency with one of the sequences designated by SEQ ID NO 1 to 15.

The present invention covers modifications of the sequences according to SEQ ID NO 1 to 15 which result e.g. from the degeneration of the genetic code, deletions, insertions, inversions and further mutations, the biological property of the encoded channel protein or part thereof being preferably maintained.

Furthermore, the invention relates to an mRNA molecule comprising a sequence corresponding to one of the above-described sequences. Accordingly the invention covers a polypeptide which is encoded by the above-mentioned nucleic acid.

The above-described sequences can be used for a screening method or also a diagnosing method. In a screening method, it is possible owing to the identification of the sequence of the I_h channel to test the effect of substances on ion channels using said sequences.

Such a screening method may e.g. comprise the following steps:

Insert 2

Insert 2:

A nucleic acid molecule as described above can be cloned into any suitable vector. The selection of vectors and methods to construct them are commonly known to persons of ordinary skill in the art and are described in general technical references (see, in general, "Recombinant DNA Part D," *Methods in Enzymology*, Vol. 153, Wu and Grossman, eds., Academic Press (1987); Birren et al., *Genome Analysis: A Laboratory Manual Series, Volume 1, Analyzing DNA*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1997); Birren et al., *Genome Analysis: A Laboratory Manual Series, Volume 2, Detecting Genes*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1998); Birren et al., *Genome Analysis: A Laboratory Manual Series, Volume 3, Cloning Systems*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1999); Birren et al., *Genome Analysis: A Laboratory Manual Series, Volume 4, Mapping Genomes*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1999); and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989)). Desirably, the vector comprises regulatory sequences, such as transcription and translation initiation and termination codons, which are specific to the type of host into which the vector is to be introduced, as appropriate and taking into consideration whether the vector is DNA or RNA. Preferably, the vector comprises regulatory sequences that are specific to the genus of the host. Most preferably, the vector comprises regulatory sequences that are specific to the species of the host.

Constructs of vectors, which are circular or linear, can be prepared to contain an entire nucleic acid sequence as described above or a portion thereof ligated to a replication system functional in a prokaryotic or eukaryotic host cell. Replication systems can be derived from ColE1, 2 μ plasmid, λ , SV40, bovine papilloma virus, and the like.

In addition to the replication system and the inserted nucleic acid, the

construct can include one or more marker genes, which allow for selection of transformed or transfected hosts. Marker genes include biocide resistance, e.g., resistance to antibiotics, heavy metals, etc., complementation in an auxotrophic host to provide prototrophy, and the like.

Suitable vectors include those designed for propagation and expansion or for expression or both. A preferred cloning vector is selected from the group consisting of the pUC, series the pBluescript series (Stratagene, LaJolla, CA), the pET series (Novagen, Madison, WI), the pGEX series (Pharmacia Biotech, Uppsala, Sweden), and the pEX series (Clonetech, Palo Alto, CA). Bacteriophage vectors, such as λ GT10, λ GT11, λ ZapII (Stratagene), λ EMBL4, and λ NM1149, also can be used. Examples of plant expression vectors include pBI101, pBI101.2, pBI101.3, pBI121 and pBIN19 (Clonetech, Palo Alto, CA). Examples of animal expression vectors include pEUK-C1, pMAM and pMAMneo (Clonetech).

An expression vector can comprise a native or nonnative promoter operably linked to an isolated or purified nucleic acid molecule as described above. The selection of promoters, e.g., strong, weak, inducible, tissue-specific and developmental-specific, is within the skill in the art. Similarly, the combining of a nucleic acid molecule as described above with a promoter is also within the skill in the art.

Thus, in view of the above, the present invention also provides a host cell comprising an isolated or purified nucleic acid molecule or a vector as described above. Examples of host cells include, but are not limited to, a human cell, a human cell line, *E. coli*, *B. subtilis*, *P. aeruginosa*, *S. cerevisiae*, and *N. crassa*. Other examples include *E. coli* TB-1, TG-2, DH5 α , XL-Blue MRF' (Stratagene), SA2821 and Y1090.

- producing homogeneous channel preparations, for example, by expression of the above-mentioned nucleic acid in a suitable host, such as oocytes, mammalian cells, etc.,
- testing of substances on said channel preparations.

It can here be determined by measuring the channel activity under the action or in the absence of test substances which substances are suited for influencing the channels.

The invention also relates to a kit for performing such a screening method which comprises at least one of the above-described nucleic acids or polypeptides.

The sequences can also be used for a diagnosing method, in particular for recognizing cardiovascular disorders.

In said diagnosing method the nucleic acid of the patient is preferably contacted with a sequence section of one of the above-described DNAs and/or RNAs, whereby a signal is obtained that is indicative of the presence and/or absence of an ion-channel nucleic acid sequence. Mutations in the ion channels of the patient can also be detected by selecting suitable samples, e.g. short oligonucleotides, which in turn is of help to the differential diagnosis.

Furthermore, the present invention refers to a kit for carrying out such a diagnosing method comprising one of the above-described sequences.

Furthermore, it is possible to use the above-described sequences for the treatment and/or prophylaxis of cardiovascular disorders and disturbances of consciousness as well as pain states. In a preferred embodiment, cardiovascular disorders that are due to

Furthermore, disturbances of consciousness that are due to a malfunction of cortico-thalamic neurons are preferably recognized. For instance, within the scope of gene

Insert 3

Insert 4

Year	Age	Sex	Weight (kg)	Height (cm)	Body mass index (kg/m ²)	Waist circumference (cm)	Waist-hip ratio	Trunk muscle strength (kg)	Trunk muscle endurance (s)	Trunk muscle power (W)
2000	20	Male	70	175	22.5	85	0.85	100	120	1200
2001	21	Female	60	160	23.4	75	0.80	90	110	1100
2002	22	Male	75	180	23.1	90	0.85	110	130	1300
2003	23	Female	65	165	23.8	80	0.82	95	115	1150
2004	24	Male	80	185	23.2	95	0.88	120	140	1400
2005	25	Female	70	170	23.9	85	0.83	105	125	1250
2006	26	Male	85	190	23.5	100	0.90	130	150	1500
2007	27	Female	75	175	24.2	90	0.86	115	135	1350
2008	28	Male	90	195	23.6	105	0.92	140	160	1600
2009	29	Female	80	180	24.7	95	0.89	125	145	1450
2010	30	Male	95	200	23.8	110	0.94	150	170	1700
2011	31	Female	85	185	24.8	100	0.91	135	155	1550
2012	32	Male	100	205	23.9	115	0.96	160	180	1800
2013	33	Female	90	190	25.1	105	0.93	145	165	1650
2014	34	Male	105	210	24.0	120	0.98	170	190	1900
2015	35	Female	95	195	25.2	110	0.95	155	175	1750
2016	36	Male	110	215	24.1	125	0.99	180	200	2000
2017	37	Female	100	200	25.0	115	0.97	165	185	1850
2018	38	Male	115	220	24.2	130	1.00	190	210	2100
2019	39	Female	105	205	25.3	120	0.99	175	195	1950
2020	40	Male	120	225	24.3	135	1.01	200	220	2200

Figure 1B shows the S4 motif of said channel protein, as compared with other known channel sequences;

Figure 1C shows the pore motif of said sequence as compared with other sequences of other channels;

Figure 1D shows the cNMP-binding domain of the cDNA of the I_h ion channel as compared with other sequences of ion channels;

Insert 3:

Accordingly, the present invention provides a method of prophylactically or therapeutically treating a mammal for a cardiovascular disorder, in particular a cardiovascular disorder that is due to a faulty control of the sinus node. The method comprises administering to a mammal (i) a vector comprising and expressing a prophylactically or therapeutically effective amount of an above-described nucleic acid or (ii) a prophylactically or therapeutically effective amount of an above-described polypeptide, whereupon the mammal is treated for the cardiovascular disorder.

The present invention further provides a method of prophylactically or therapeutically treating a mammal for a disturbance of consciousness, in particular a disturbance of consciousness that is due to a malfunction in thalamic neurons. The method comprises administering to a mammal (i) a vector comprising and expressing a prophylactically or therapeutically effective amount of an above-described nucleic acid or (ii) a prophylactically or therapeutically effective amount of an above-described polypeptide, whereupon the mammal is treated for the disturbance of consciousness.

Still further provided by the present invention is a method of prophylactically or therapeutically treating a mammal for a pain state. The method comprises administering to a mammal (i) a vector comprising and expressing a prophylactically or therapeutically effective amount of an above-described nucleic acid or (ii) a prophylactically or therapeutically effective amount of an above-described polypeptide, whereupon the mammal is treated for the pain state.

002180-22504950

Insert 4:

Therefore, the present invention also provides a composition comprising an above-described isolated or purified nucleic acid (or vector comprising the nucleic acid) or an above-described polypeptide and a carrier therefor. Carriers, such as pharmaceutically acceptable carriers, are well-known in the art, and are readily available. The choice of carrier will be determined in part by the particular route of administration and whether a nucleic acid molecule or a polypeptide molecule is being administered. Accordingly, there is a wide variety of suitable formulations for use in the context of the present invention, and the invention expressly provide a pharmaceutical composition that comprises an active agent of the invention and a pharmaceutically acceptable carrier therefor. The following methods and carriers are merely exemplary and are in no way limiting.

Formulations suitable for oral administration can consist of (a) liquid solutions, such as an effective amount of the compound dissolved in diluent, such as water, saline, or orange juice; (b) capsules, sachets or tablets, each containing a predetermined amount of the active ingredient, as solids or granules; (c) suspensions in an appropriate liquid; and (d) suitable emulsions. Tablet forms can include one or more of lactose, mannitol, corn starch, potato starch, microcrystalline cellulose, acacia, gelatin, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, stearic acid, and other excipients, colorants, diluents, buffering agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible excipients. Lozenge forms can comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth. Pastilles can comprise the active ingredient in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like containing, in addition to the active ingredient, such excipients/carriers as are known in the art.

Formulations suitable for parenteral administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain anti-oxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The

09640582 081700

formulations can be presented in unit-dose or multi-dose sealed containers, such as ampules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid excipient, for example, water, for injections, immediately prior to use. Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described. Further suitable formulations are found in Remington's Pharmaceutical Sciences, 17th ed., (Mack Publishing Company, Philadelphia, Pa.: 1985), and methods of drug delivery are reviewed in, for example, Langer, Science, 249, 1527-1533 (1990).

Generally, when an above-described polypeptide is administered to an animal, such as a mammal, in particular a human, it is preferable that the polypeptide is administered in a dose of from about 1 to about 1,000 micrograms of the polypeptide per kg of the body weight of the host per day when given parenterally. However, this dosage range is merely preferred, and higher or lower doses may be chosen in appropriate circumstances. For instance, the actual dose and schedule can vary depending on whether the composition is administered in combination with other pharmaceutical compositions, or depending on interindividual differences in pharmacokinetics, drug disposition, and metabolism. One skilled in the art easily can make any necessary adjustments in accordance with the necessities of the particular situation.

If desired, the half-life of the polypeptide can be increased by conjugation to soluble macromolecules, such as polysaccharides, or synthetic polymers, such as polyethylene glycol, as described, for instance, in U.S. Patents 5,116,964, 5,336,603, and 5,428,130. Alternately, the polypeptides can be "protected" in vesicles composed of substances such as proteins, lipids (for example, liposomes), carbohydrates, or synthetic polymers. If liposomes are employed, liposome delivery can be carried out as described in U.S. Patent 5,468,481, or using liposomes having increased transfer capacity and/or reduced toxicity *in vivo* (see, e.g., PCT patent application WO 95/21259 and the references cited therein). Furthermore, polypeptides can be administered in conjunction with adenovirus (preferably replication-deficient adenovirus) to allow the intracellular uptake of the polypeptides by adenoviral-mediated uptake of bystander molecules (e.g.,

007F80 28507950

as described in PCT patent application WO 95/21259). Similarly, a conjugate, such as one comprising a targeting moiety, or a fusion of an above-described polypeptide to an antibody (or an antigenically reactive fragment thereof) that recognizes a cell surface antigen; etc. can be employed to deliver the resultant fusion protein to a specific target cell or tissue (e.g., as described in U.S. Patent 5,314,995).

Those of ordinary skill in the art can easily make a determination of the vector to be administered to an animal, such as a mammal, in particular a human. The dosage will depend upon the particular method of administration, including any vector or promoter utilized. For purposes of considering the dose in terms of particle units (pu), also referred to as viral particles, it can be assumed that there are 100 particles/pfu (e.g., 1×10^{12} pfu is equivalent to 1×10^{14} pu). An amount of recombinant virus, recombinant DNA vector or RNA genome sufficient to achieve a tissue concentration of about 10^2 to about 10^{12} particles per ml is preferred, especially of about 10^6 to about 10^{10} particles per ml. In certain applications, multiple daily doses are preferred. Moreover, the number of doses will vary depending on the means of delivery and the particular recombinant virus, recombinant DNA vector or RNA genome administered.

Further provided by the present invention is a hybridoma cell line that produces a monoclonal antibody that is specific for an above-described isolated or purified polypeptide molecule. Methods of making hybridomas are known in the art (see, e.g., Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1988); Harlow et al., *Using Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1999)). Thus, the present invention also provides the monoclonal antibody produced by the hybridoma cell line. Similarly, the present invention provides a polyclonal antiserum raised against an above-described isolated or purified polypeptide molecule. Methods of raising polyclonal antiserum against a polypeptide molecule are also known in the art (see, e.g., Harlow et al. (1988), *supra*; Harlow et al. (1999), *supra*).

0078012250450

Figure 2A shows the inward current having a complex waveform, which is triggered by the hyperpolarizing voltage steps from a holding voltage of +10 mV to more negative test values;

Figure 2B shows the equilibrium current/voltage (I/V) relationship determined at the end of a hyperpolarizing voltage pulse;

Figure 2C shows the measuring protocol for the determination of the "instantaneous" I/V relationship from the amplitude of the tail currents;

Figure 2D shows the "instantaneous" I/V relationship which is slightly outwardly rectifying, at a reversal voltage V_{rev} of -30 mV;

Figure 2E shows that the time course of the "tail" currents depends on the time of the change in voltage;

Figure 2F shows the voltage dependence of the relative probability that the channel is open, P_o , which was determined from the amplitude of tail currents at +10 mV, similar to those illustrated in Fig. 2A;

Figure 3A shows the induction of large whole-cell currents by hyperpolarization in the presence of 1 mM cAMP, which currents developed with a delay and slowly reached an equilibrium;

Figure 3B shows the voltage dependence of P_o , determined from normalized whole-cell "tail" currents and "tail" currents of inside-out patches;

Figure 4G shows that the inward currents were interrupted almost entirely, whereas the amplitudes of the outward currents did not change when the extracellular medium just contained Na^+ ;

Figure 4H shows the I/V relationship of the currents from part G at different K^+ concentrations;

Figure 5A shows a Northern Blot of the channel messenger RNA with a major transcript of about 3.3 kb and a minor transcript of 2.9 kb;

Figure 5B is a light-microscopic photograph of sperms from *S. purpuratus* (right picture) and the corresponding immunohistochemical staining with an antibody which specifically recognizes the SPIH channel (left picture).

Figure 5C shows a corresponding Western Blot analysis.

Figure 6 is a schematic illustration showing the pc SPIH construct that was used for the heterologous expression of SPIH in HEK 293 cells. The cDNA region is illustrated as a hatched bar; the adjoining regions of the plasmid vector (pcDNA I) as bold lines. The orientation of the cDNA in the plasmid vector can be inferred from the position of the promoter for the T7 polymerase and the restriction sites in the multiple cloning region. The inserted Kozak sequence is designated by K.

A typical representative of an ion channel protein according to the invention is the channel from sea urchin (SPIH). The channel activity of HEK 293 cells, which had been transfected with the pcSPIH construct (Fig. 6), was examined with the help of the patch-clamp method in the whole-cell configuration. Hyperpolarizing voltage steps showed an inward current with a complex waveform (cf. Fig. 2A). A fast current component that was not time-resolved was followed by a time-dependent current that developed with a delay and, after the maximum had been reached, decreased into smaller amplitudes when the test voltage was $V_m \leq -30$ mV (Fig. 2A). After V_m had been set back to +10 mV, "tail" currents developed that also showed a complex time course. The steady-state

relationship between current/voltage (I/V), at the end of the hyperpolarizing voltage pulse (arrow in Fig. 2A), showed a strong inward rectification (Fig. 2B). The "instantaneous" I/V relationship was determined from the amplitude of the tail currents using a different protocol for the voltage steps (Fig. 2C). The "instantaneous" I/V relationship was slightly outwardly rectifying with a reversal voltage, V_{rev} , of -30 mV (Fig. 2D). The I/V relationship became approximately linear at higher $[K^+]_o$ because the inward sodium current was significantly amplified by $[K^+]_o$ (see Fig. 4H). The conclusion can be drawn that the currents are strongly inwardly rectifying because the SPIH channel at positive voltages is either closed or inactivated. The voltage dependence of the open probability, P_o (Fig. 2F), was determined from the amplitude of the tail currents at $+10$ mV (Fig. 2A). The voltage, $V_{1/2}$, at which a half-maximal current was observed, was at -26.1 mV (7 experiments). Thus the SPIH channel is inactive at voltages $\geq +10$ mV and is opened by hyperpolarization. This voltage dependence reminds of hyperpolarization-activated currents (I_h) which occur in different cells (DiFrancesco, 1990, 1993; Pape, 1996). Because of its unusual properties, the I_h has also been designated as a "queer" or "funny" current (I_q and I_f). The channel according to the invention is (1) activated at hyperpolarizing voltages; (2) directly modulated by cyclic nucleotides; (3) blocked by millimolar concentrations of extracellular Cs^+ ; (4) it is cation-selective at a P_{Na}/P_K of ~ 0.2 to 0.4 ; and (5) the inward sodium currents are sensitive to $[K^+]_o$. The following experiments demonstrate that said features are also found in the heterologously expressed SPIH channel.

With 1 mM cAMP in the pipette solution, hyperpolarization produced large currents which developed with a delay and slowly reached a steady state (Fig. 3A). The sigmoidal time course of the current (see Fig. 3A, box) is characteristic of the time course of vertebrate I_h currents. 1 mM cGMP in the pipette also changed the SPIH-induced currents. The voltage dependence of P_o was determined with the help of whole-cell tail currents (Fig. 3B). A fit to the Boltzmann equation yielded $V_{1/2} = -50.8$ mV. The dialysis of the cell with

the pipette solution took several minutes; thus transient effects of cAMP might impair the test. A technique was therefore employed using the rapid photorelease of cAMP or cGMP from "caged" derivatives (cf. Adams and Tsien, 1993; Hagen et al., 1996). The cells were dialyzed with 100 μ M "caged" cAMP and the SPIH channels were activated by changing the V_m from +10 mV to -70mV; a short flash of UV light then effected a rapid increase in the amplitude of the SPIH-induced inward current (Fig. 3C). The hyperpolarization-activated currents before the flash resembled control currents (Fig. 3C, trace 1), while amplitude and time course of the currents after the UV flash (Fig. 3C, trace 2) were similar to those recorded in the presence of cAMP (Fig. 3E). With 100 μ M "caged" cGMP in the pipette, UV flashes of similar duration and similar intensity did not change the SPIH-induced currents. A binding motif for cyclic nucleotides suggests that cAMP could directly enhance the channel activity without the participation of a phosphorylation mechanism. To verify this hypothesis the SPIH currents were measured on excised membrane patches without (Fig. 3D) and with cAMP (Fig. 3E). cAMP (1mM) enhanced the amplitudes of the voltage-activated currents by up to 20-fold. The increase in current by cAMP was reversible and did not require Mg^{2+} /ATP. The superfusion of the excised membrane patches with solutions containing different cAMP concentrations enhanced the SPIH currents in a dose-dependent way. The dependence of the current on the cAMP concentration can be described by a simple binding isotherm with a $K_{1/2}$ of 0.74 μ M and a Hill coefficient which does not significantly differ from one (Fig. 3F). In the separated membrane patches, $V_{1/2}$ in the presence of cAMP was about 35 mV more negative than $V_{1/2}$ measured in the whole-cell configuration (Fig. 3B). This observation might suggest that an endogenous factor provided by the HEK293 cell also determines $V_{1/2}$. cGMP concentrations of up to 1mM did not change the amplitude of the SPIH currents. The conclusion can be drawn from this experiment that cAMP, but not cGMP, can modulate the SPIH channel activity. Thus, in contrast to CNG channels (Finn et al., 1996) SPIH is under the double control of voltage and cAMP. Blockage of the SPIH channels by extracellular Cs^+ was examined on "outside-out" membranes with the

voltage protocol of Fig. 2C. Cs^+ blocked the SPIH channel in a concentration- and voltage-dependent manner. In the presence of 10 mM Cs^+ the inward currents disappeared completely, whereas outward tail currents were still present (cf. Figs. 4A and 4B). The I/V relationship in the presence of from 0 to 10 mM Cs^+ is shown in Fig. 4C. The standardized current I/I_{\max} (at -70 mV) was plotted against $[\text{Cs}^+]$ (Fig. 4D). The data were fitted with an inhibitory constant K_i of 245 μM and a Hill coefficient of $n = 1.2$. The ion selectivity of the SPIH channel was determined with inside-out membranes. The bath solutions always contained 0.1 mM cAMP to increase the amplitude of the currents. 100 mM K^+ in the bath were replaced by Rb^+ , Na^+ , Li^+ or Cs^+ (Fig. 4E). The permeability ratios $P_{\text{K}} : P_{\text{Rb}} : P_{\text{Na}} : P_{\text{Li}} : P_{\text{Cs}}$ were calculated as 1 : 0.7 : 0.26 : 0.15 : 0.06. The ion selectivity of SPIH concurs well with the ion selectivity of various vertebrate I_{h} channels (Pape, 1996; Wollmuth and Hille, 1992). When the extracellular medium only contained Na^+ , the inward currents were eliminated almost entirely, whereas the amplitudes of the outward currents did not change significantly (Fig. 4G). Elevation of $[\text{K}^+]_o$ to 5 and 20 mM dramatically increased the inward currents. These results demonstrate that the SPIH channel conducts little, if any, sodium in the absence of potassium ions.

The expression of messenger RNA of the channel protein was analyzed by means of Northern Blots. A major transcript of around 3.3 kb and a minor transcript of 2.9 kb were detected in poly(A)⁺RNA of male, but not female, gonads (Fig. 5A). The size of the transcripts concurs well with the size of the cloned cDNA (3 kb). The SPIH-specific probe did not hybridize with poly(A)⁺RNA isolated from the intestine of sea urchin (Fig. 5A). The exclusive expression of SPIH mRNA in male gonads suggests that the channel is expressed in sperms. This hypothesis was tested with purified antibodies FPc44K and FPc45K directed against a fusion protein of the C-terminal domain of the channel polypeptide (residues 662-767). The antibodies were used for Western Blot analyses (Fig. 5C) and immunocytochemistry (Fig. 5B). Both antibodies recognized a main band of $M_r \sim 92\text{K}$ in Western Blots of flagellar membranes which had been purified from sea

urchin sperm (Fig. 5C, lane 3). Membranes which had been purified from the sperm head were not recognized by the antibodies (Fig. 5C, lane 5). This result was confirmed by immunocytochemistry with individual sperms. The antibody FPc45K almost exclusively stained the sperm flagellum (Fig. 5B); the weak staining of some head structures presumably represents unspecific cross reactivity of the antibody. A band of $M_r \sim 88K$ was observed in Western Blots of membranes of transfected HEK293 cells (Fig. 5C, lane 2). The M_r of the channel polypeptide, expressed in HEK293 cells, is almost identical with the M_r value as is to be expected of the derived amino acid sequence (87.9K). In membranes of non-transfected HEK293 cells, no 88K polypeptide was detected by the antibody (Fig. 5C, lane 1). The treatment of flagellar membranes with alkaline phosphatase lowered the M_r of the native polypeptide from $\sim 92 K$ to 88K. Since native and heterologously expressed polypeptides were of a similar size, the cloned cDNA carries the complete coding sequence of SPIH. The small decrease in M_r under dephosphorylating conditions demonstrates that the native polypeptide in phosphorylated form is present with a slightly reduced electrophoretic mobility. In most dephosphorylation experiments the shift from 92K to 88K was not complete, and at least two intermediate bands were observed. This result suggests that the channel polypeptide should be phosphorylated several times. The SPIH sequence carries sequence motifs for the phosphorylation by PKA, PKG, PKC and tyrosine kinase (see Fig. 1A).

The electrophysiological properties unequivocally identify SPIH as a member of the I_h channel family. However, we also noticed characteristic differences between SPIH and vertebrate I_h channels. First, in the absence of cAMP the SPIH current is transient, whereas in the presence of cAMP the time course is similar to that in vertebrate I_h channels. Second, the large augmentation of the SPIH current by cAMP primarily arises from an increase in the maximum current while cAMP modulates the cardiac I_h channel such that $V_{1/2}$ is shifted towards more positive values (DiFrancesco, 1993) without

influencing the maximum amplitudes (see, however, Ingram and Williams, 1996; Accili et al., 1997). Finally, the cardiac I_h is also modulated by micromolar cGMP concentrations (DiFrancesco and Tortora, 1991), whereas SPIH does not exhibit said effect. The SPIH channel is very similar to both the voltage-controlled K^+ channels and the CNG cation channels. That is why the I_h channels form a class of their own within the superfamily of the voltage-controlled channels. SPIH has a characteristic motif of a voltage sensor (S4) like the K^+ , Na^+ and Ca^{2+} channels that are opened by depolarization. Although there is no a priori reason to rule out the S4 motif as a voltage sensor in a hyperpolarization-activated channel, the mechanism of an activation as in HERG- K^+ channels (Trudeau et al., 1995; Smith et al., 1996) is more likely. It has been demonstrated with respect to the strong inward rectification of HERG that it is the result of the inactivation which closes the channels at positive voltages, but the channels recover rapidly from the inactivation at negative voltages. In HERG channels the inactivation is much faster than the activation and is therefore not visible kinetically (Smith et al., 1996). Together with the CNG channels SPIH possesses a cyclic nucleotide-binding region, and its properties are modulated by cAMP. cAMP probably intensifies the SPIH activity by binding to the highly conserved cyclic nucleotide-binding region. In CNG channels, it has been demonstrated with respect to the high selectivity for cGMP that said selectivity is accompanied by a Thr residue (T363 in the α -subunit of the rod photoreceptor; Altenhofen et al., 1991) and an Asp residue (D604 in rCNG α ; Varnum et al., 1995). The SPIH has Val and Ile residues at the corresponding positions; it is presumed that these positions also control the ligand selectivity in SPIH. The physiological importance of the I_h channels in flagellar membranes of sperm could be explained as follows: the stimulation of *S. purpuratus* sperm with the chemotactic peptide "speract" causes a hyperpolarization (Lee and Garbers, 1986; Garbers, 1989), of which it is assumed that it is due to the opening of a K^+ channel (Babcock et al., 1992). At higher peptide concentrations the hyperpolarization is followed by a depolarization (Babcock et al., 1992). Two (or more) ion channel types with different selectivity and pharmacology could contribute to the "speract"-induced

002180 22504950

depolarization (see Darszon et al, 1996). One of said channels has a weak K^+ selectivity ($P_{Na/PK} \cong 0.2$) and an extremely low P_o (at $V_m = 0$ mV) which is considerably enhanced by cAMP, but not by cGMP (Labarca et al., 1996). These observations suggest that said channel is actually SPIH. The "speract"-induced hyperpolarization could initiate the SPIH channel activity which then could even be augmented by a simultaneous increase in the cAMP level (Hansbrough et al., 1980) with the help of a voltage-dependent adenylate cyclase (Beltrán et al, 1996). At the given ionic composition of sea water and a $P_{Na/PK}$ of 0.2 to 0.4 the opening of the SPIH channel and the subsequent Na^+ influx could effect the "speract"-induced depolarization. It can also reasonably be assumed that the I_h channels, for instance in cardiac cells or thalamic neurons, take part in the generation of oscillations of the membrane voltage, thereby causing the oscillation of Ca^{2+} in the flagellum (Suarez et al, 1993). The change in $[Ca^{2+}]_i$, could change the flagellar beating, thereby contributing to the chemotactic response.

0040582-081700

EXAMPLES

Methods

Isolation of the cDNA clones

With two degenerated primers (# 1764 and # 1772) a PCR was carried out on single-strand cDNA (from sea urchin gonads, *Drosophila melanogaster*, bovine retina, olfactory tissue of the rat) or on cDNA libraries (from human thalamus or heart). A 100 µl PCR batch had the following composition: 3-10 ng of first-strand cDNA and about 10⁵ pfu of the cDNA libraries, respectively, 1.6 µg of the degenerated primer each, 1 x PCR buffer, 2mM dNTP, 1 U PrimeZyme (Biometra). The PCR batch was first denatured at 94°C for 2 min and then incubated for 45 cycles in the following manner:

denaturation:	94°C, 45 sec
hybridization:	48°C, 45 sec
polymerization:	72°C, 40 sec

The sequences of the degenerated primers are (in 5'→3' direction):

1764: CTGACTGCAGARGTNTTYCARCCNGGNGA (SEQ ID NO 16)

1772: ATCGGAATTCNCCRAARTANGANCCRTC (SEQ ID NO 17)

The PCR fragments amplified with the primers # 1764 and # 1772 were radiolabeled and used as probes for screening cDNA libraries under high stringency for the complete cDNAs. The partial clone HHIH (SEQ ID NO 11) was isolated by low-stringency hybridization. The hybridization conditions were as follows:

	high stringency	low stringency
prehybridization	5 x SSC ⁽¹⁾ , 5 x Denhardt's ⁽²⁾ , 0.1% SDS, 0.1 mg/ml herring sperm DNA, 1-2 h, 65°C	5 x SSC ⁽¹⁾ , 5 x Denhardt's ⁽²⁾ , 0.1% SDS, 0.1 mg/ml herring sperm DNA, 1-2 h, 55°C
hybridization	prehybridization solution with 50-100 ng ³² P-labeled DNA (1- 10 ⁶ cpm/ml), 12-14 h, 65°C	prehybridization solution with 50-100 ng ³² P-labeled DNA (1x10 ⁶ cpm/ml), 12-14 h, 65°C
washing	1 x SSC(1), 0.1% SDS 2 x 30 min, 65°C	2 x SSC(1), 0.1% SDS 2 x 30 min, 55°C

⁽¹⁾ 1 x SSC 150 mM NaCl, 15 mM Na citrate, pH 7.0

⁽²⁾ 1 x Denhardt's Ficoll, polyvinylpyrrolidone, bovine serum albumin (0.2 g/l each)

The positive phages were isolated and the cDNA was converted by "in vivo excision" (in case of λ ZAPII phages) into pBluescriptSK derivatives. The cDNA was excised with EcoRI from λ gt11 phages and subcloned into pBluescriptSK plasmid DNA. The DNA was sequenced with the dideoxy-mediated chain termination technique (Sanger et al., 1997).

Northern and Western Blots

Poly(A)⁺RNA, isolated from different sea urchin tissues, was analyzed by Northern blotting. Each lane contained about 10 μ g poly(A)⁺RNA. The blot was hybridized with a ³²P-labeled 1074 bp cDNA fragment (nucleotide positions) at 42°C, 5 x SSC and 50% formamide. A C-terminal region of the SPIH polypeptide was expressed as a fusion construct with the maltose binding protein. The purified fusion protein was used for producing the polyclonal antibodies FPc44K and FPc45K; the antibodies were purified from rabbit serum by affinity chromatography using the fusion protein. Sperm flagella were separated from the head according to Darszon et al. (1994). Purified flagella and

head membranes were homogenized in a solution buffer containing 150 mM NaCl, 1 mM $MgCl_2$, 20 mM Hepes at pH 7.5, 0.1 mM EGTA and 0.5% Triton X-100, followed by a centrifugation at 40,000 rpm for 60 minutes. This process was repeated two times. Transfected HEK293 cells were homogenized in a lysis buffer (10 mM Hepes, 1 mM DTT and 1 mM EDTA at pH 7.4), 5 x freeze-dried (in liquid N_2) and finally centrifuged at 55,000 rpm for 10 minutes. The membrane pellet was dissolved in the solution buffer. Flagellar membrane proteins were dephosphorylated with a unit of alkaline phosphatase in solution buffer at 30°C for 30 to 60 min. The membrane proteins were separated by SDS-PAGE, transferred to Immobilon membranes and labeled with the polyclonal antibodies. The immunoreactivity was made visible by the ECL detection kit (Amersham). Immunocytochemistry on an individual sperm was carried out as described above (Weiner 1997).

Electrophysiology

cDNA coding the SPIH polypeptide was transiently expressed in HEK293 cells, as described earlier (Baumann et al, 1994). SPIH-controlled currents were recorded with the patch-clamp method in the whole-cell configuration and cell-free membrane patches. The composition of various bath and pipette solutions is indicated in the legends of the figures (see below). The channels were activated by stepping the membrane voltage from +10 mV to various negative voltage values. Leakage currents were subtracted using a P/8 protocol. The voltage dependence of the probability that the channel is open was determined from tail currents at +10 mV. The blockade of the SPIH channel by Cs^+ was analyzed with outside-out membrane patches in the presence of 1 mM cAPM in a pipette solution. The solutions in the bath contained 0.03 to 10 mM CsCl. Relative ion permeabilities were calculated from the respective shift of V_{rev} , which was measured on cell-free inside-out membrane patches, when 100 mM K^+ in the bath had been replaced

by Na^+ , Li^+ , Rb^+ or Cs^+ . Experiments with "caged" cAMP or "caged" cGMP were carried out as described earlier (Hagen et al. 1996).

The results of said experiments are now described in more detail.

Figure 1A shows the nucleic acid sequence and the derived amino acid sequence of the I_h channel of sea urchin (SPIH). Nucleotides are numbered in 5'→3' direction, +1 corresponding to the first nucleotide of the start codon (ATG) of the open reading frame. Nucleotides that are 5'-located from nucleotide +1 are designated by negative figures. The derived amino acid sequence (one-letter code) is indicated under the nucleic acid sequence and is also numbered. The start codon (ATG), the corresponding methionine and the stop codon (TGA; pos. 2302-2304) are printed in bold. Stop codons in the same reading frame before the start codon are underlined. The polyadenylation signal at position 2501-2507 is boxed. The position of the transmembranal segments S1-S6, of the pore-forming region and of the binding site for cyclic nucleotides (cNMP binding site) is marked by bars above the nucleic acid sequence. The limits of said regions are defined by sequence comparison with other voltage-dependent K^+ channels, EAG- K^+ channels and CNG channels. Consensus sequences for phosphorylation by cAMP/cGMP-dependent kinases are marked by triangles (Δ). Consensus sequences for phosphorylation by protein kinase C are marked by circles (\bullet) and that by tyrosine kinase by an asterisk (\ast). The SPIH sequence (SEQ ID NO 4) codes for a protein of 767 amino acids with a calculated molecular weight of 87,937 Da.

Figure 1B shows a comparison of the voltage-sensor (S4) motifs of the I_h channel of sea urchin and other channels. Regularly spaced Arg or Lys residues are boxed. Other positively charged residues are in bold.

Shaker (Pongs et al., 1988), K^+ channel encoded by the *Drosophila* Shaker gene;

DmEAG (Warmke et al, 1997), *Drosophila* EAG channel;
 HERG, human EAG-related gene (Warmke and Ganetzky, 1994);
 KAT1 (Anderson et al, 1992), K⁺ channel of *Arabidopsis thaliana*;
 brCNGC α (Kaupp et al, 1989), α -subunit of the cyclic nucleotide-controlled channel from bovine rod photoreceptors.

Figure 1C shows the pore motif of SPIH with the pore motifs of other members of the superfamily of the voltage- and cyclic nucleotide-controlled ion channels:

The residues which are identical or similar to the corresponding amino acids in SPIH are highlighted by a black or grey background.

Figure 1D shows a sequence comparison of cNMP binding domains.

boCNGC α , the α -subunit of the CNG channel of bovine olfactory neurons (Ludwig et al., 1990); PKA1, the cAMP binding site 1 of the protein kinase A (Titani et al., 1984); the cGMP binding site 1 of the protein kinase G (Takio et al, 1984); CAP, the catabolite activator protein (Aiba et al., 1982). Residues that are highly conserved in cyclic nucleotide-binding motifs are indicated by arrows; residues that determine the ligand selectivity in brCNGC α are indicated by an asterisk. Secondary-structure predictions derived from the cAMP binding domain of CAP are shown as bars below the sequence.

Figure 2 shows the electrophysiological characterization of the SPIH channel.

Figure 2A shows the current, which was recorded by transfected HEK293 cells in the whole-cell configuration. The current was activated by stepping the voltage from a holding value at +10 mV to various test values of -100 mV to +10 mV in increments of 10 mV. Tail currents were recorded by stepping the voltage of the test value back to +10 mV. The HEK293 cells were flushed with a bath solution containing the following (mM):

135 NaCl, 5 KCl, 1.8 CaCl₂, 2.8 MgCl₂ and 5 Hepes-NaOH at pH 7.4; the pipette solution contained the following substances (mM): 126 KCl, 10 Hepes-KOH, 10 EGTA at pH 7.4.

In **Figure 2B**, there is plotted the voltage-current (I/V) relationship measured under equilibrium conditions at the time indicated by the arrowhead in Figure 2A.

Figure 2C shows the measurement protocol with which the "instantaneous" I/V relationship was determined; the voltage was first stepped from a holding value of 0 mV to -70 mV, followed by steps to test values in the range of from +50 mV to -70 mV in 10 mV increments.

Figure 2D then shows the plot of the "instantaneous" I/V relationship measured at the time indicated by the arrow in Figure 2C (inset).

Figure 2E shows that the time course of the "tail" currents depends on the time at which the voltage is reset to + 30 mV.

Figure 2F shows the voltage dependence of the relative open probability, P_o , of the channel. The tail current amplitudes (arrow in part a) were normalized to the maximum current. The midpoint voltage, $V_{1/2}$, was -26.1 mV. The effective charge amount, Q , which is flowing during channel switching, is 3.5 elementary charges. It was achieved from a fit of the Boltzmann function to the data: Mean of 7 experiments.

Figure 3 indicates the modulation of SPIH channels by cyclic nucleotides.

Figure 3A shows the whole-cell SPIH current in the presence of 1 mM cAMP. The voltage-step protocol is the same as in Fig. 2A. The bath contained (mM): 135 NaCl, 5 KCl, 1.8 CaCl₂, 2.8 MgCl₂ and 5 Hepes-NaOH at pH 7.4; the pipette solution contained

(mM): 126 KCl, 10 Hepes-KOH, 10 EGTA at pH 7.4, and 1mM cAMP. The inset shows a magnification by way of which the sigmoidal time course can be seen particularly well.

Figure 3B shows the voltage dependence of the relative P_o , derived from normalized whole-cell tail currents at +10mV (●) and of tail currents recorded by inside-out patches (Δ). A continuous line represents a fit of the Boltzmann equation to the data. $V_{1/2}$ for the whole-cell currents of part A was -50.8mV and for the inside-out-patch currents of part E it was -84.7 mV; the Q values were 3.8 and 2.7, respectively.

Figure 3C shows the modulation of whole-cell SPIH currents by the photolysis of "caged" cAMP. The pipette solution contained 100 μ M "caged" cAMP. The SPIH current was activated by voltage jumps from +10 mV to -70 mV before the UV flash was induced (trace 1) and after three consecutive UV flashes (trace 2). The time course of the flash-induced increase in current at -70 mV is shown below.

Figures 3D and E show voltage-activated SPIH currents in inside-out membrane patches without cAMP (D) and in the presence of 1 mM cAMP (E) in the bath. The voltage step protocol was carried out in the way as shown in Figure 2A. The pipettes and bath solutions contained (mM): 126 KCl, 10 Hepes-KOH, 10 EGTA at pH 7.4 and 1 mM cAMP (bath).

Figure 3F discloses the dependence of the SPIH current amplitude on the cAMP concentration; the cAMP concentrations were as follows (μ M): 0.1; 0.3; 1; 3; 10 and 1000. A continuous line shows a fit of the Hill equation to the data; $K_{1/2} = 0.74 \mu$ M; $n = 1.05$; mean of 10 experiments.

Figure 4 shows several pharmacological properties of the SPIH channel.

Figures 4A and B show voltage-activated SPIH currents, recorded by outside-out membrane patches without (A) and with 10 mM Cs⁺ (B) in the bath; the pipette solution contained the following (mM): 124 KCl, 10 Hepes-KOH, 10 EGTA at pH 7.4 and 1 mM cAMP; the bath solution contained (mM): 126 KCl, 10 Hepes-KOH, 10 EGTA at pH 7.4 and the illustrated concentrations of CsCl.

Figure 4C shows again the I/V relationship in the presence of 0 to 10 mM Cs⁺ in the bath.

Figure 4D discloses the dependence of the normalized current at -70 mV on [Cs⁺]. The continuous line shows a fit of the Hill equation to said data; K_i = 245 μM, Hill coefficient 1.2 (mean of 1-6 experiments).

Figure 4E shows the ion selectivity of the SPIH channel. V_{rev} was determined on inside-out patches by stepping the holding voltage (-70 mV) to test values between -30 mV and +30 mV in 5 mV increments. The pipette solution contained the following (mM): 150 KCl, 10 Hepes-NMDG, 10 EGTA at pH 7.4; the bath solution was composed as follows (mM): 50 KCl, 100 XCl, 10 Hepes-NMDG, 10 EGTA at pH 7.4 and 0.1 cAMP.

Figure 4F shows the I/V relationship of the currents shown in part E. V_{rev} was 16.9 mV (Na⁺), 20.6 mV (Li⁺), 5.6 mV (Rb⁺), and 24.6 mV (Cs⁺); mean of 3 to 10 experiments. The relative ion permeabilities P_X/P_K were calculated according to the equation $P_X/P_K = \{[K^+]_o - [K^+]_i \exp(zFV_{rev}/RT)\} / [X^+]_i \exp(zFV_{rev}/RT)$.

Figure 4G shows the K⁺ dependence of whole-cell inward Na⁺ currents in the presence of 0.5 mM and 20 mM K⁺ in extracellular medium.

Figure 4H shows the "instantaneous" I/V relationship in the presence of 0, 1, 3, 5, 10, and 20 mM K^+ in the bath.

The pipette solution was the same as in part B, the bath solution as in Figure 1A with the indicated K^+ concentrations; the ion intensities were adjusted to the same value by the respective NMDG concentrations.

Figure 5 shows the expression pattern of SPIH.

Figure 5A is a Northern Blot analysis of the tissue distribution of SPIH transcripts in mRNA of male gonads (lane 1), female gonads (lane 2) and intestinal cells (lane 3); 10 μ l poly(A)⁺RNA each.

Figure 5B is a Western Blot analysis of membranes of mock-transfected HEK293 cells (lane 1; 2.5 μ g protein), HEK293 cells which were transfected with SPIH cDNA (lane 2; 2.5 μ g protein), purified flagella from sperm of *S. purpuratus* (lane 3; 6 μ g protein), dephosphorylated flagellar membranes (lane 4; 6 μ g protein) and sperm heads (lane 5; 15 μ g protein).

Reference table of the DNA sequences described in the text by SEQ ID numbers

SEQ ID NO	DNA sequence
1	Partial sequence of the I_h channel from human thalamus tissue
2	partial sequence of an I_h channel from olfactory rat tissue
3	partial sequence of an I_h channel from retinal bovine tissue
4	complete sequence of the I_h channel from sea urchin sperm
5	complete sequence of the I_h channel from <i>Drosophila melanogaster</i>
6	partial sequence of an I_h channel from retinal bovine tissue
7	partial sequence of an I_h channel from retinal bovine tissue
8	partial sequence of an I_h channel from olfactory rat tissue
9	partial sequence of an I_h channel from olfactory rat tissue
10	partial sequence of an I_h channel from human thalamus tissue
11	partial sequence of an I_h channel from human heart tissue
12	complete sequence of an I_h channel from retinal bovine tissue
13	partial sequence of an I_h channel from olfactory rat tissue
14	partial sequence of an I_h channel from olfactory rat tissue
15	complete sequence of an I_h channel from human heart tissue

Literature

Accili, E.A., Redaelli, G. and DiFrancesco, D.: Differential control of the hyperpolarization-activated current (i) by cAMP gating and phosphatase inhibition in rabbit sino-atrial node myocytes. *J.Physiol.* 500 (1997) 643-651.

Adams, S.R. and Tsien, R.Y.: Controlling cell chemistry with caged compounds. *Annu.Rev.Physiol.* 55 (1993) 755-784.

Aiba, H., Fujimoto, S. and Ozaki, N.: Molecular cloning and nucleotide sequencing of the gene for *E. coli* cAMP receptor protein. *Nucleic Acids Res.* 10 (1982) 1345-1361.

Altenhofen, W., Ludwig, J., Eismann, E., Kraus, W., Bönigk, W. and Kaupp, U.B.: Control of ligand specificity in cyclic nucleotide-gated channels from rod photoreceptors and olfactory epithelium. *Proc.Natl.Acad.Sci.USA* 88 (1991) 9868-9872.

Anderson, J.A., Huprikar, S.S., Kochian, L.V., Lucas, W.J. and Gaber, R.F.: Functional expression of a probable *Arabidopsis thaliana* potassium channel in *Saccharomyces cerevisiae*. *Proc.Natl.Acad.Sci.USA* 89 (1992) 3736-3740.

Araki, T., Ito, M. and Oshima, T.: Potential changes produced by application of current steps in motoneurons. *Nature* 191 (1962) 1104-1105.

Attwell, D. and Wilson, M.: Behavior of the rod network in the tiger salamander retina mediated by membrane properties of individual rods. *J.Physiol.* 309 (1980) 287-315.

Babcock, D.F., Bosma, M.M., Battaglia, D.E. and Darszon, A.: Early persistent activation of sperm K channels by the egg peptide speract. *Proc.Natl.Acad.Sci.USA* 89 (1992) 6001-6005.

Bader, C.R., MacLeish, P.R. and Schwartz, E.A.: A voltage-clamp study of the light response in solitary rods of the tiger salamander. *J.Physiol.* 296 (1979) 1-26.

Bader, C.R., Bertrand, D. and Schwartz, E.A.: Voltage-activated and calcium-activated currents studied in solitary rod inner segments from the salamander retina. *J.Physiol.* 331 (1982) 253-284.

Baumann, A., Frings, S., Godde, M., Seifert, R. and Kaupp, U.B.: Primary structure and functional expression of a *Drosophila* cyclic nucleotide-gated channel present in eyes and antennae. *EMBO J.* 13 (1994) 5040-5050.

Beltrán, C., Zapata, O. and Darszon, A.: Membrane potential regulates sea urchin sperm adenylylcyclase. *Biochemistry* 35 (1996) 7591-7598.

Brown, H.F. and DiFrancesco, D.: Voltage clamp investigations of current underlying pacemaker activity in rabbit-sino-atrial node. *J. Physiol.* 308 (1980) 221-251.

Brown, H.F., DiFrancesco, D. and Noble, S.J.: How does adrenaline accelerate the heart? *Nature* 280 (1979) 235-236.

Darszon, A., Labarca, P., Beltrán, C., García-Soto, J. and Liévano, A.: Sea urchin sperm: An ion channel reconstitution study case. *Methods: A Companion to Methods in Enzymology* 6 (1994) 37-50.

Darszon, A., Liévano, A. and Beltrán, C.: Ion channels: Key elements in gamete signaling. In *Current Topics in Developmental Biology*, Vol. 44. Academic Press, San Diego, 1996, pp.117-167.

Dayhoff, M.O., Schwartz, R. M., Orcutt, B.C. (1978) in: *Atlas of protein sequence and structure*, Band 5, Suppl. 3, Hrsg.: Dayhoff, M.O., National Biomedical Research Foundation, Silver Spring, MD, S. 345-352.

DiFrancesco, D.: A new interpretation of the pace-maker current in calf Purkinje fibres. *J. Physiol.* 314 (1981a) 359-376.

DiFrancesco, D.: A study of the ionic nature of the pace-maker current in calf Purkinje fibres. *J. Physiol.* 314 (1981b) 277-293.

DiFrancesco, D.: The hyperpolarization-activated current, i_h , and cardiac pacemaking. In Rosen, M.R., Janse, M.J. and Wit, A.L. (Eds.), *Cardiac Electrophysiology: a Textbook*. Futura, New York, 1990, pp.117-132.

DiFrancesco, D.: Pacemaker mechanisms in cardiac tissue. *Annu.Rev.Physiol.* 55 (1993) 455-472.

DiFrancesco, D. and Tortora, P.: Direct activation of cardiac pacemaker channels by intracellular cyclic AMP. *Nature* 351 (1991) 145-147.

Fain, G.L., Quandt, F.N., Bastian, B.L. and Gerschenfeld, H.M.: Contribution of a caesium-sensitive conductance increase to the rod photoresponse. *Nature* 272 (1978) 467-469.

Finn, J.T., Grunwald, M.E. and Yau, K.-W.: Cyclic nucleotide-gated ion channels: An extended family with diverse functions. *Annu.Rev.Physiol.* 58 (1996) 395-426.

Garbers, D.L.: Molecular basis of fertilization. *Annu.Rev.Biochem.* 58 (1989) 719-742.

Garbers, D.L.: Guanylyl cyclase receptors and their endocrine, paracrine, and autocrine ligands. *Cell* 71 (1992) 1-4.

Hagen, V., Dzeja, C., Frings, S., Bendig, J., Krause, E. and Kaupp, U.B.: Caged compounds of hydrolysis-resistant analogues of cAMP and cGMP: Synthesis and application to cyclic nucleotide-gated channels. *Biochemistry* 35 (1996) 7762-7771.

Halliwel, J.V. and Adams, P.R.: Voltage-clamp analysis of muscarinic excitation in hippocampal neurons. *Brain Res.* 250 (1982) 71-92.

Hansbrough, J.R., Kopf, G.S. and Garbers, D.L.: The stimulation of sperm metabolism by a factor associated with eggs and by 8-bromo-guanosine 3',5'-monophosphate. *Biochim.Biophys.Acta* 630 (1980) 82-91.

Hille, B.: Ionic channels of excitable membranes. Sinauer Associates Inc., Sunderland, 1992.

Hodgkin, A.L. and Huxley, A.F.: A quantitative description of membrane current and its application to conduction and excitation in nerve. *J.Physiol.* 117 (1952) 500-544.

Ingram, S.L. and Williams, J.T.: Modulation of the hyperpolarization-activated current (I) by cyclic nucleotides in guinea-pig primary afferent neurons. *J.Physiol.* 492 (1996) 97-106.

Ito, M. and Oshima, T.: Electrical behavior of the motoneurone membrane during intracellularly applied current steps. *J. Physiol.* 180 (1965) 607-635.

Kaupp, U.B., Niidome, T., Tanabe, T., Terada, S., Bönigk, W., Stühmer, W., Cook, N.J., Kangawa, K., Matsuo, H., Hirose, T., Miyata, T. and Numa, S.: Primary structure and functional expression from complementary DNA of the rod photoreceptor cyclic GMP-gated channel. *Nature* 342 (1989) 762-766.

Labarca, P., Santi, C., Zapata, O., Morales, E., Beltrán, C., Liévano, A. and Darszon, A.: A cAMP regulated K-selective channel from the sea urchin sperm plasma membrane. *Develop.Biol.* 174 (1996) 271-280.

Lee, H.C. and Garbers, D.L.: Modulation of the voltage-sensitive Na/H exchange in sea urchin spermatozoa through membrane potential changes induced by the egg peptide speract. *J.Biol.Chem.* 261 (1986) 16026-16032.

Llinás, R.R.: The intrinsic electrophysiological properties of mammalian neurons; insights into central nervous system function. *Science* 242 (1988) 1654-1664.

Ludwig, J., Margalit, T., Eismann, E., Lancet, D. and Kaupp, U.B.: Primary structure of cAMP-gated channel from bovine olfactory epithelium. *FEBS Lett.* 270 (1990) 24-29.

Pape, H.-C.: Queer current and pacemaker: The hyperpolarization-activated cation current in neurons. *Annu.Rev.Physiol.* 58 (1996) 299-327.

Pongs, O., Kecskemethy, N., Müller, R., Krah-Jentgens, I., Baumann, A., Kiltz, H.H., Canal, I., Llamazares, S. and Ferrus, A.: *Shaker* encodes a family of putative potassium channel proteins in the nervous system of *Drosophila*. *EMBO J.* 7 (1988) 1087-1096.

Sanger, F., Nicklen, S. and Coulson, A.R.: DNA sequencing with chain-terminating inhibitors. *Proc.Natl.Acad.Sci.USA* 74 (1977) 5463-5467.

Smith, P.L., Baukrowitz, T. and Yellen, G.: The inward rectification mechanism of the HERG cardiac potassium channel. *Nature* 379 (1996) 833-836.

Suarez, S.S., Varosi, S.M. and Dai, X.: Intracellular calcium increases with hyperactivation in intact, moving hamster sperm and oscillates with the flagellar beat cycle. *Proc.Natl.Acad.Sci.USA* 90 (1993) 4660-4664.

Takio, K., Wade, R.D., Smith, S.B., Krebs, E.G., Walsh, K.A. and Titani, K.: Guanosine cyclic 3',5'-phosphate dependent protein kinase, a chimeric protein homologous with two separate protein families. *Biochemistry* 23 (1984) 4207-4218.

Titani, K., Sasagawa, T., Ericsson, L.H., Kumar, S., Smith, S.B., Krebs, E.G. and Walsh, K.A.: Amino acid sequence of the regulatory subunit of bovine type I adenosine cyclic 3',5'-phosphate dependent protein kinase. *Biochemistry* 23 (1984) 4193-4199.

Trudeau, M.C., Warmke, J.W., Ganetzky, B. and Robertson, G.A.: HERG, a human inward rectifier in the voltage-gated potassium channel family. *Science* 269 (1995) 92-95.

Varnum, M.D., Black, K.D. and Zagotta, W.N.: Molecular mechanism for ligand discrimination of cyclic nucleotide-gated channels. *Neuron* 15 (1995) 619-625.

Warmke, J., Drysdale, R. and Ganetzky, B.: A distinct potassium channel polypeptide encoded by the *Drosophila eag* locus. *Science* 252 (1991) 1560-1562.

Warmke, J.W. and Ganetzky, B.: A family of potassium channel genes related to *eag* in *Drosophila* and mammals. *Proc.Natl.Acad.Sci.USA* 91 (1994) 3438-3442.

Weiner, J.: Molekularbiologische, immunologische und funktionelle Charakterisierung von β -Untereinheiten des zyklisch Nukleotid-gesteuerten Ionenkanals aus dem Rinderhoden. Dissertation (1996) Universität Düsseldorf

Wollmuth, L.P. and Hille, B.: Ionic selectivity of I_h channels of rod photoreceptors in tiger salamanders. *J.Gen.Physiol.* 100 (1992) 749-765.

Yanagihara, K. and Irisawa, H.: Inward current activated during hyperpolarization in the rabbit sino atrial node cell. *Pflügers Arch.* 385 (1980) 11-19.

WHAT IS CLAIMED IS:

1. An isolated or purified nucleic acid which codes for an I_h ion channel or a part thereof, or a nucleic acid complementary thereto, except for the nucleic acids with the sequences indicated in the GenBank database under the accession number AF028737 and in the ENHUM database under the accession number N72770, wherein said nucleic acid is DNA or RNA.
2. The isolated or purified nucleic acid of claim 1, characterized in that the nucleic acid is of human origin.
3. The isolated or purified nucleic acid of claim 2, characterized in that the nucleic acid comprises the sequence according to SEQ ID NO 1 or a part thereof.
4. The isolated or purified nucleic acid of claim 1, characterized in that the nucleic acid is of rat origin.
5. The isolated or purified nucleic acid of claim 4, characterized in that the nucleic acid comprises the sequence according to SEQ ID NO 2 or a part thereof.
6. The isolated or purified nucleic acid of claim 1, characterized in that the nucleic acid is of bovine origin.
7. The isolated or purified nucleic acid of claim 6, characterized in that the nucleic acid comprises a sequence according to SEQ ID NO 3 or SEQ ID NO 12 or a part of either of the foregoing sequences.
8. The isolated or purified nucleic acid of claim 1, characterized in that the nucleic acid is of sea urchin origin.
9. The isolated or purified nucleic acid of claim 8, characterized in that the nucleic acid comprises the sequence according to SEQ ID NO 4 or a part thereof.
10. The isolated or purified nucleic acid of claim 1, characterized in that the nucleic acid is of *Drosophila* origin.

002780-28504950

11. The isolated or purified nucleic acid of claim 10, characterized in that the nucleic acid comprises the sequence according to SEQ ID NO 5 or a part thereof.

12. An isolated or purified nucleic acid characterized in that the sequence thereof is at least 80% identical to the isolated or purified nucleic acid of SEQ ID NO 1, 2, 3, 4 or 12.

13. The isolated or purified nucleic acid of claim 12, characterized in that the sequence thereof is at least 90% identical to the isolated or purified nucleic acid of SEQ ID NO 1, 2, 3, 4 or 12.

14. An isolated or purified nucleic acid characterized in that the nucleic acid hybridizes under low stringency conditions with SEQ ID NO 1, 2, 3, 4, 5 and/or 12.

15. The isolated or purified nucleic acid of claim 14, characterized in that the nucleic acid hybridizes under stringent conditions with SEQ ID NO 1, 2, 3, 4, 5 and/or 12.

16. A vector comprising the isolated or purified nucleic acid of claim 1.

17. A host cell comprising the vector of claim 16.

18. A composition comprising the isolated or purified nucleic acid of claim 1 and a carrier therefor.

19. An isolated or purified polypeptide encoded by a nucleic acid of claim 1.

20. An isolated or purified polypeptide encoded by a nucleic acid of claim 3.

21. An isolated or purified polypeptide encoded by a nucleic acid of claim 5.

22. An isolated or purified polypeptide encoded by a nucleic acid of claim 7.

23. An isolated or purified polypeptide encoded by a nucleic acid of claim 9.

24. An isolated or purified polypeptide encoded by a nucleic acid of claim 11.

002720 "22504350

25. An isolated or purified polypeptide encoded by a nucleic acid of claim 12.
26. An isolated or purified polypeptide encoded by a nucleic acid of claim 13.
27. An isolated or purified polypeptide encoded by a nucleic acid of claim 14.
28. An isolated or purified polypeptide encoded by a nucleic acid of claim 15.
29. A composition comprising the isolated or purified polypeptide of claim 19 and a carrier therefor.
30. A monoclonal antibody that specifically binds to the polypeptide of claim 19.
31. A method of screening a substance for the ability to influence the activity of an I_h ion channel, which method comprises:
 - (a) providing a homogeneous I_h ion channel preparation,
 - (b) contacting the homogeneous I_h ion channel preparation with the substance,
 - (c) measuring the activity of the I_h ion channel preparation in the presence of the substance, and
 - (d) comparing the activity of the I_h ion channel preparation in the presence of the substance with the activity of the I_h ion channel preparation in the absence of the substance, wherein a change in the activity of the I_h ion channel preparation in the presence of the substance as compared to the activity of the I_h ion channel preparation in the absence of the substance indicates that the substance can influence the I_h ion channel.
32. The method of claim 31, wherein said I_h ion channel preparation is prepared by expressing the isolated or purified nucleic acid which codes for an I_h ion channel or a part thereof, or a nucleic acid complementary thereto, except for the nucleic acids with the sequences indicated in the GenBank database under the accession number AF028737 and in the ENHUM database under the accession number N72770, wherein said nucleic acid is DNA or RNA, in a host cell.
33. The method of claim 31, wherein said I_h ion channel preparation consists essentially of the polypeptide encoded by the isolated or purified nucleic acid which codes for an I_h ion channel or a part thereof, or a nucleic acid complementary thereto, except for the

004052-00400

nucleic acids with the sequences indicated in the GenBank database under the accession number AF028737 and in the ENHUM database under the accession number N72770.

34. A method of diagnosing an I_h ion channel-associated disorder in a patient, which method comprises:

- (a) contacting a nucleic acid sample from said patient with a detectably labeled isolated or purified nucleic acid of claim 1 under hybridizing conditions,
- (b) detecting the label of the detectably labeled isolated or purified nucleic acid molecule, and
- (c) comparing the level of detection of the label in (b) with the level of detection of the label in a control sample, wherein a difference in the level of detection of the label in (b) and the level of detection of the label in a control sample is indicative of an I_h ion channel-associated disorder in a patient.

35. The method of claim 34, wherein the detectably labeled isolated and purified nucleic acid is mutated, in which case the detection of the label in (b) is indicative of the presence of a nucleic acid encoding a mutated I_h ion channel in the nucleic acid sample of the patient.

36. The method of claim 34, wherein said I_h ion channel-associated disorder is a cardiovascular disorder.

37. A method of prophylactically or therapeutically treating a mammal for a cardiovascular disorder, which method comprises administering to said mammal a vector comprising and expressing a prophylactically or therapeutically effective amount of an isolated or purified nucleic acid of claim 1, whereupon said mammal is treated for said cardiovascular disorder.

38. The method of claim 37, wherein said cardiovascular disorder is due to a faulty control of the sinus node.

39. A method of prophylactically or therapeutically treating a mammal for a cardiovascular disorder, which method comprises administering to said mammal a prophylactically or therapeutically effective amount of a polypeptide of claim 19, whereupon said mammal is treated for said cardiovascular disorder.

004200-22504950

40. The method of claim 39, wherein said cardiovascular disorder is due to a faulty control of the sinus node.

41. A method of prophylactically or therapeutically treating a mammal for a disturbance of consciousness, which method comprises administering to said mammal a vector comprising and expressing a prophylactically or therapeutically effective amount of an isolated or purified nucleic acid of claim 1, whereupon said mammal is treated for said disturbance of consciousness.

42. The method of claim 41, wherein said disturbance of consciousness is due to a malfunction in thalamic neurons.

43. A method of prophylactically or therapeutically treating a mammal for a disturbance of consciousness, which method comprises administering to said mammal a prophylactically or therapeutically effective amount of a polypeptide of claim 19, whereupon said mammal is treated for said disturbance of consciousness.

44. The method of claim 43, wherein said disturbance of consciousness is due to a malfunction in thalamic neurons.

45. A method of prophylactically or therapeutically treating a mammal for a pain state, which method comprises administering to said mammal a vector comprising and expressing a prophylactically or therapeutically effective amount of an isolated or purified nucleic acid of claim 1, whereupon said mammal is treated for said pain state.

46. A method of prophylactically or therapeutically treating a mammal for a pain state, which method comprises administering to said mammal a prophylactically or therapeutically effective amount of a polypeptide of claim 19, whereupon said mammal is treated for said pain state.

09640521 081700

Abstract

Sequences of an I_h ion channel and use thereof

The present invention relates to a nucleic acid, preferably a DNA, comprising at least a part of the sequence of an I_h ion channel. Said sequence may e.g. be derived from a human DNA, a rat DNA, a bovine DNA, a *Drosophila melanogaster* DNA or a sea urchin DNA. Furthermore, the present invention relates to an mRNA molecule which contains the corresponding sequences. The invention further relates to a polypeptide or protein encoded by the nucleic acid.

Furthermore, the invention relates to the use of the inventive nucleic acid and proteins for a screening and/or diagnosing method and to the kits required therefor.

Lastly, the invention relates to the use of one or more nucleic acids and proteins for the treatment and/or prophylaxis of cardiovascular disorders and disturbances of consciousness.

Sequence ID No. 4

CGGGAGAATAGTGCACCAAGGGATGCCCGTGA	AAATATTAATTAACAGTTTTTTAAGAACAA	-101
TCATCAAACCCGGGCCCCATCATGAAGGAATAACAAGGCCTTCGAAAAGTATGGGAAACT		-41
GGTCGGCAGGACATCAGCATTATTAATTCTAGGAAACTCATTATGGATAACAAGGAAACT		18
	M D N K E T	6
AACGGAGAGCTAGAGCAGTCTGATGAGGCCGATCCGTCCGGTCAAAACCTTGATGATGGG		78
N G E L E Q S D E A D P S G Q N L D D G		26
GAAACCGATAGCAAACAAGAAGAGAATCTCATCAACGTTAGCCCGCCAAAAACACCGCCA		138
E T D S K Q E E N L I N V S P P K T P P		46
GGTCCTCCTCCTCCTCTAAAGAATGGAGGAAGGGGTCAGAAACCGCCCAAAATCCCAATA		198
G P P P P L K N G G R G Q K P P K I P I		66
TGTCATCAAAATGGAAAGCTCCCCAAGGAAGTTGAATGGACAGAAGACAGAGGCGAAGAC		258
C H Q N G K L P K E V E W T E D R G E D		86
AGAAAGGATAGTCTCACTCTTCAATCAAAGCTAGATCACGGGGCATACACGGATGAGAAA		318
R K D S L T L Q S K L D H G A Y T D E K		106
	▲	
CAGGATCTTCTAACATATCTTGACCGTCACGGCATCAACAGTCCAGTCAAGCTAACACCA		378
Q D L L T Y L D R H G I N S P V K L T P		126
GATGAAACTGGAGGGAGCAGTGCTTTGGATATTCTTGGGATTATTGAAGAGAGGGGACACT		438
D E T G G S S A L D I L G I I E E R D T		146
GGTGCCTAGGCTCTGATCCCTCATCCACTATGCAGGCCATGGCTAAACCTGTAGGCTTT		498
G A L G S D P S S T M Q A M A K P V G F		166
CTGCAGAGGCAGCTATGGACTGTCTCTCCAACCTTCAGACAATAGACTCTCCATGAAACTT		558
L Q R Q L W T V L Q P S D N R L S M K L		186
	●	
TTCGGAAGCAAGAAAGGGTTACAAAAGGAAAAATATCGGCTGAGGAAGGCGGGGGTTCTT		618
F G S K K G L Q K E K Y R L R K A G V L		206
	●	
	S1	
ATCATTTCATCCATGTAGTCATTTTCAGATTTTACTGGGATCTACTGATGCTGTGCCTGATC		678
I I H P C S H F R F Y W D L L M L C L I		226

ATGGCAAACGTCATCCTCCTACCCGTCGTCATTACTTTCTTCCACAACAAGGACATGAGT		738
M A N V I L L P V V I T F F H N K D M S		246

	S2	
ACGGGTTGGCTCATCTTTAATTGCTTCTCAGATACCTTCTTCATTCTCGATCTCATCTGC		798
T G W L I F N C F S D T F F I L D L I C		266

AACTTTTCGGACCGGCATCATGAATCCGAAGTCGGCCGAACAGGTGATCCTCAACCCCCGT		858
N F R T G I M N P K S A E Q V I L N P R		286

	S3	
CAAATCGCCTATCATTATCTCCGTTTCATGGTTCATCATCGATCTCGTGTCTTCCATC		918
Q I A Y H Y L R S W F I I D L V S S I P		306

ATGGACTACATCTTCCTCCTCGCTGGCGGCCAGAACCGTCACTTCCTCGAGGTGTCCCGA 978
 M D Y I F L L A G G Q N R H F L E V S R 326

S4

GCCCTCAAGATACTGCGCTTTGCCAAGCTCCTCAGTCTTCTTCGACTCCTGCGTCTGTCC 1038
 A L K I L R F A K L L S L L R L L R L S 346

AGGCTCATGCGGTTCGTCAGTCAATGGGAACAGGCCTTCAACGTAGCCAATGCCGTCATC 1098
 R L M R F V S Q W E Q A F N V A N A V I 366

S5

CGGATCTGTAATCTAGTGTGTATGATGCTTCTGATTGGCCATTGGAATGGCTGCCTTCAA 1158
 R I C N L V C M M L L I G H W N G C L Q 386

TATCTCGTGGCCATGCTGCAAGAATACCCCGACCAATCATGGGTGCGCCATTAATGGCCTT 1218
 Y L V P M L Q E Y P D Q S W V A I N G L 406

Pore

GAGCAGCTCATTGGTGGGAGCAGTATACATGGGCACTCTTCAAAGCCCTTTCCGACATG 1278
 E H A H W W E Q Y T W A L F K A L S H M 426

CTCTGTATCGGGTACGGCAAGTTCCCCCCTCAAAGCATCACCGATGTCTGGCTAACGATT 1338
 L C I G Y G K F P P Q S I T D V W L T I 446

S6

GTCAGTATGGTGTCCGGTGGCGACCTGCTTCGCCCTGTTTCATCGGACACGCTACCAATCTC 1398
 V S M V S G A T C F A L F I G H A T N L 466

ATCCAGTCCATGGACTCCTCCAGCAGGCAATACCGTGAGAAGTTGAAACAAGTTGAAGAG 1458
 I Q S M D S S S R Q Y R E K L K Q V E E 486

TACATGCAGTATCGCAAGCTACCGTCCCACCTACGAAACAAGATCCTCGATTACTACGAG 1518
 Y M Q Y R K L P S H L R N K I L D Y Y E 506

TACCGATACCGAGGAAAGATGTTTGATGAGAGGCATATCTTTCGAGAAGTGTCGGAGAGT 1578
 Y R Y R G K M F D E R H I F R E V S E S 526

ATACGACAGGATGTCGCAAACTACAATTGTGCGGACCTGGTGCATCCGTCCCTTTCTTC 1638
 I R Q D V A N Y N C R D L V A S V P F F 546

GTCGGTGCCGACTCAAACCTTCGTCACCCGTGTGGTGACGCTGCTCGAATTCGAGGTCTTC 1698
 V G A D S N F V T R V V T L L E F E V F 566

CAACCCGCTGACTATGTTATACAGGAAGGTACTTTTCGGTGATCGCATGTTCTTCATCCAG 1758
 Q P A D Y V I Q E G T F G D R M F F I Q 586

CAGGGCATCGTCGACATCATCATGTCCGACGGCGTCATCGCCACGTCACTCAGTGACGGC 1818
 Q G I V D I I M S D G V I A T S L S D G 606

cNMP binding site

TCATATTTTGGCGAAATCTGCCTGCTTACCCGTGAGCGCCGCGTGGCATCGGTGAAGTGC 1878
 S Y F G E I C L L T R E R R V A S V K C 626

GAGACCTACTGCACGCTCTTCTCGCTCTCCGTCCAGCATTTCAACCAAGTGCTCGACGAG 1938
 E T Y C T L F S L S V Q H F N Q V L D E 646

TTTCCCGCCATGAGGAAAACGATGGAAGAGATAGCCGTTTCGTCTGACCCGAATCGGG 1998
 F P A M R K T M E E I A V R R L T R I G 666
 AAGGAATCGAGCAAGCTGAAATCCCGCCTAGAGAGCCCGACGATCAGGGACACTGCCCCCT 2058
 K E S S K L K S R L E S P T I R D T A P 686
 CTCCTTCCGATCCACCTGATACACCGTCTTTTCGTACCCGACATCGAAAAGAACCGGTTTC 2118
 L F P I P P D T P S F V T D I E K N R F 706
 TTTGGCGACGACACGGACGATGTACACATCAGGACCCGAGTCGACGTCGAGCGTGGTTCCG 2173
 F G D D T D D V H I R T R V D V E R G S 726
 CATGAAAACGTCATCGCCATCATGGATGGGAGTTTATCCGACCTCAGGATGGAAAACGAA 2238
 H E N V I A I M D G S L S D L R M E N E 746
 ATCCAAGCCCGTAAATCGTCTAGCGGAAAACGGAGGAAATTCACGCAACAAACACCGAA 2293
 I Q A R K S S S G K R R K F Q Q Q T T E 756
 CTATGACGACTTGAAACAAACAATGATGGACGCTTACAATTTCCAGTGATTCAATACTTA 2358
 L - 767
 CGCAATGCAGACATTAGCTTTTGTACCTGATTGTTTAGAATGTATTGAATTTGTAGATCA 2413
 GTCCCGGCAATAAGAAAGCATAATTTGGAATTTCTTTTCATTGAGGAAGTACTGAAAACAA 2478
 TGTGATAGCAGCCCGGTAGAAATTTCTTGTCCATTATCGAGGCTATATTTTTTCGCGCTTTC 2538
 TTACGAAGTAAATGAAAGGATCAATTAATTTATTGTTCTTTGTCTCCGTGCGCTTTGTATC 2598
 TGATGCCGAAAAGGAATGAAACGTGATTAGAACAGTAATCGATTGAACTACAGAAAGTCTT 2658
 TTCAAAATGTTGAATGTATGAAGGAGGAGGGGGAAGGTTTGATATATGCAAAGAAATGGA 2713
 GAAATATTTTTGTAAATTTATCTAGAATGGTACTATTGATGCTGGAAGGTGTTGAAGTT 2778
 GTCCAATATTGTGTCAAATCACCAACTATTTGACATTTGTCTTTTTC 2825

0940563 081700

Fig. 1B, 1C

	S4 motif																pore																																				
B																																	C																				
SPIH	326-	R	A	K	I	L	R	F	A	K	L	L	S	L	L	R	L	L	R	L	S	R	L	M	R	-350	416-	T	W	A	L	F	K	A	E	S	H	M	L	C	I	G	Y	G	K	F	P	P	Q	S	-438		
Shaker	344-	M	S	L	A	I	L	R	V	I	R	L	V	R	V	F	R	I	F	K	L	S	R	H	S	K	-368	418-	P	D	A	F	W	W	A	V	V	T	M	T	T	V	G	Y	G	D	M	T	P	V	G	+40	
DmEAG	341-	S	L	F	S	A	L	K	V	V	R	L	L	R	L	G	R	V	V	R	K	L	D	R	Y	L	-365	441-	V	T	A	L	V	F	T	M	T	C	M	T	S	V	G	F	G	N	V	A	A	R	T	-463	
HERG	519-	E	L	I	G	L	L	K	T	A	R	L	L	R	L	V	R	V	A	R	K	L	D	R	Y	S	-343	612-	V	T	A	L	Y	F	T	F	S	L	T	S	V	G	F	G	N	V	S	P	N	T	-634		
KAT I	168-	S	M	L	R	L	W	R	L	R	R	V	S	S	L	F	A	R	L	E	K	D	I	R	F	N	-192	248-	V	T	A	L	V	W	S	T	T	L	T	T	T	T	G	Y	G	D	F	H	A	E	N	-270	
brCNGC α	263-	W	N	Y	P	E	I	R	L	N	R	L	L	R	I	S	R	M	F	E	F	F	Q	R	T	E	-287	348-	V	T	S	L	T	W	S	T	L	T	L	T	T	T	I	G	-	-	E	T	P	P	P	V	-368

Fig. 1D

cNMP binding domain

	553-FOUR	485-L	462-L	579-C	750-C	143-E	110-Q	10-T	αA	$\beta 1$	$\beta 2$	$\beta 3$	$\beta 4$	$\beta 5$	$\beta 6$																								
SPIH	VU	VL	LE	FE	VF	QF	DY	VI	QEG	T	FG	DR	MF	FI	Q	GG	I	VD	II	NS	D	--	GV	A	--	TS	LS	DG	SV	FG	E	I							
brCNGC α	VE	VL	KL	LP	VL	SP	DY	CK	KG	D	IG	RE	VI	IK	KG	KH	AV	AD	--	--	--	GT	Q	F	V	VL	SD	GS	YF	GE	I								
boCNGC α	VE	VL	KL	LP	QV	SP	DY	CR	KG	D	IG	KE	VI	IK	KG	KL	AV	AD	--	--	--	GV	T	Q	Y	A	L	LS	AG	SC	FG	E	I						
DmEAG	CR	AM	H	F	M	S	AP	DL	YH	T	GE	S	DS	CF	I	V	T	GS	FE	VI	Q	DD	--	EV	A	--	I	L	SK	D	V	FG	Q						
HERG	CR	AM	K	F	K	T	HA	PG	TH	HA	G	DL	TA	VI	F	IS	R	GS	FE	II	R	SD	--	V	V	A	--	I	L	E	K	ND	I	F	GE	P			
PKA I	CR	DF	AM	F	P	V	SF	IA	ST	TV	IQ	GD	EG	DN	FI	VI	D	Q	GE	D	Y	UN	E	--	W	A	--	TS	V	GE	GG	SG	FG	E	I				
PKG I	Q	Q	ES	DT	F	DC	M	Y	P	VE	G	K	D	SC	IK	EG	D	V	GS	LV	V	MD	GV	EV	TK	--	GV	K	--	LC	MG	PG	KV	FG	E	I			
CAP	TE	WF	ES	CH	I	H	K	Y	PS	K	S	T	I	H	G	GE	K	ET	RY	IV	K	GS	V	AV	DI	K	D	EE	G	K	EM	IL	SY	L	N	GD	FI	GE	I

SPIH	C L L	- - - -	R R R V A S	V K C E T Y C T L F S L S	V O H F N Q V L D E P P A M R K T M E E I A V R R L T P I G R E S S	-670
bCNGC α	S L L N I K G S K A G	R R T A N N	K S I G Y S D L F C L S K F	D L M E A L T E P P D A N G M L E E K G K Q I L M K D G L L D I	-609	
bCNGC α	S L L N I K G S K M G	R R T A N N	L R S L G Y S D L F C L S K F	D L M E A L T E P P D A N G M L E E K G K Q I L M K D G L L D I	-586	
DmEAG	F W K D S	- - - -	V G Q S A A N V R A L T Y C D L H H A	K R D Y L L E V L D F P A F A N S F A R N L V L T Y N L R H R L I F	-697	
HERG	L N L Y A R	- - - -	P G S N G D V R A L T Y C D L H K H	R R D L L E V L D M P E F S D H F W S S L E I T F N L R D T N M I	-868	
PKA I	A L L Y	- - - -	G T P R A A V V K A K T N V K L Y G	D R D S V R R L L M G S L R R R K M Y E F L S K V S I L E S L D	-258	
PKG I	A L L Y N	- - - -	C T R T A T V K K T L V N V K L W A H	D R Q C F O T T M M R T G L L K H T E Y M E F L S V P T F Q S L P E	-226	
CAP	G L F F E E	- - - -	G O E R S A W V R A K T A C E V A E I S Y K K F R Q H Q V N P D L M R R S A Q M A R R L Q V T S E K V G	-132		

↓★

★


αB

β8


β7

αC

Fig. 2A, 2B



Fig, 2C, 2D



Fig, 2E, 2F




Fig. 3A, 3B





Fig. 3C, 3D





Fig. 3E, 3F

E



F




Fig. 4




Fig. 4F




Fig. 4G




Fig. 4H




Fig. 5





Fig. 6



SEQUENCE LISTING

<110> Forschungszentrum Jülich GmbH

<120> Sequences of an I_h ion channel and use thereof

<130> PCT981

<140> PCT/EP99/00942

<141> 1999-02-12

<150> DE 198 06 581.7

<151> 1998-02-17

<160> 18

<170> PatentIn Ver. 2.1

<210> 1

<211> 1342

<212> DNA

<213> Homo sapiens

<400> 1

```

cgttgcgctt  caccaagatc  ctcagcctcc  tgcggctgct  gcgcctctca  cgctgatcc  60
gctacatcca  tcagtgggag  gagatcttcc  acatgacctg  tgacctggcc  agcgcggtga  120
tgaggatctg  caatctcatc  agcatgatgc  tgctgctctg  ccactgggac  ggctgcctgc  180
agttcctggt  gcccatgctg  caggacttcc  cgcgcaactg  ctgggtgtcc  atcaatggca  240
tggtgaacca  ctcgtggagt  gaactgtact  ccttcgcact  cttcaaggcc  atgagccaca  300
tgctgtgcat  cgggtacggc  cggcagggcg  ccgagagcat  gacggacatc  tggctgacca  360
tgctcagcat  gattgtgggt  gccacctgct  acgcatgtt  catcgccac  gccactgcc  420
tcattcagtc  gctggactcc  tcgcggcgcc  agtaccagga  gaagtacaag  caggtggagc  480
agtacatgtc  cttccacaag  ctgccagctg  acttcgcca  gaagatccac  gactactatg  540
agcaccgtta  ccagggaag  atgtttgacg  aggacagcat  cctgggcgag  ctcaacgggc  600
ccctgcggga  ggagatcgtc  aacttcaact  gccggaagct  ggtggcctcc  atgccgctgt  660
tcgccaacgc  cgacccaac  ttcgtaacgg  ccagtctgac  caagctcaag  ttcgaggtct  720
tccagccggg  tgactacatc  atccgcgaag  gcaccatcgg  gaagaagatg  tacttcatcc  780
agcacggcgt  ggtcagcgtg  ctactaagg  gcaacaagga  gatgaagctg  tccgatggct  840
cctacttcgg  ggagatctgc  ctgctcacc  ggggcgcgcg  cacggcgagc  gtgctggctg  900
acacctactg  ccgcctctat  tcgctgagcg  tggacaactt  caacgaggtg  ctggaggagt  960
accccatgat  gcggcgcgcc  ttcgagacgg  tggccatcga  ccgcctggac  cgcacggca  1020
agaagaattc  catcctcctg  cacaagggtg  agcatgacct  caactcgggc  gtattcaaca  1080
accaggagaa  cgccatcatc  caggagatcg  tcaagtacga  ccgcgagatg  gtgcagcagg  1140
ccgagctggg  ctcagcgcg  gggcctcttc  ccgcgcgcgc  cgccgcgcgc  gcagtcacct  1200
cggccatcgc  cacgctgcag  caggcgcgcg  ccagtgcctt  ctgcccgcag  tggcgcggcc  1260
gctcgtgggg  ccgctggcgc  tcggctcgcc  gcgcctcgtg  cgchgcyndy  hcccggggsc  1320
cgcacctgch  gccnccac  cc

```

1342

<210> 2

<211> 3112

<212> DNA

<213> Rattus rattus

<400> 2

```

cctggttcgt  ggtggacttc  atctcctcga  tcccgggtgga  ttatatcttt  cttattgtag  60
agaaaggaat  ggattcggaa  gtttacaaga  ccgccagagc  acttcggatc  gtgaggttta  120
caaaaattct  cagtctcttg  cgtttattac  gcctttcaag  gtttaattaga  tacatacacc  180
agtgggaaga  gatattccac  atgacatatg  atctcgccag  tgcagtgggtg  agaattctca  240
acctcattgg  catgatgctg  ctctgtgtgc  actgggatgg  ctgtcttcag  tttctggctc  300

```

09640580-081700

ccttgcctgca ggacttccca ccggaattgct ggggtttctct aaatgaaatg gttaatgatt 360
 catgggggaa acagtattcc tacgcactct tcaaagctat gagtcacatg ctgtgcattg 420
 gttatggcgc ccaggccccc gtcagcatgt ctgacctctg gattaccatg ctgagcatga 480
 ttgttggggc cacctgctat gccatgtttg tccggccatgc cacagctttg atccagtctc 540
 tggattcttc aaggaggcag tatcaagaga agtacaagca agtagagcaa tacatgtcat 600
 tccacaagtt accagctgac atgcgccaga agatacatga ttactatgag caccgatacc 660
 aaggcaagat cttcgatgag gaaaatattc tcagtgaact taatgatcct ctgagagagg 720
 aaatagtcaa cttcaactgc cggaaactgg tggccaccat gcctctcttt gctaaccgag 780
 atcccaattt cgtgacggcc atgctgagca agctgagatt tgagggtgtc cagcccggag 840
 actatatcat tcgagaagga gctgtgggga agaaaatgta tttcatccag catgggtgtg 900
 ctgggtgtcat caccaagtcc agtaaagaaa tgaagttgac agacggctct tactttggag 960
 aaatatgcct gctgaccaag ggccggcgca ctgccagtgt tcgagctgat acatactgtc 1020
 gcctttactc cttttcggtg gacaatttca acgaggtctt ggaggaatat ccaatgatga 1080
 gaagagcctt tgagacagtt gctattgacc gactagatcg gataggcaag aaaaactcta 1140
 ttctcctgca gaagttccag aaggatctga aactgggtgt tttcaacaac caggagaatg 1200
 agatcctgaa gcagattgtg aagcatgaca gagagatggt acaagcgatc cctccaatca 1260
 actatcctca aatgacagcc ctgaattgca catcttcaac caccacccca acgtcgcgca 1320
 tgaggaccca atctccacca gtctacacag cgaccagcct ctctcacagc aacctgcact 1380
 caccagccc cagcacacag acgcctcaac cctcagccat cctttcaccg tgctcctaca 1440
 ccacgcagt ctgcagctct cctatacaga gcccctggc cacgcgaact ttccattatg 1500
 cctctcccac tgcattccaa ttgtcactca tgcagcagcc tcagccgcag ctacagcaat 1560
 cccaggtaca gcagactcag ccgagcgcgc agccgcagcc gcagcagccg caacagcaac 1620
 aacagcagca acagcagcag cagcagcagc agcaacaaca acagcagcag caacagccac 1680
 agacacctgg tagttccaca ccgaaaaatg aagtgcacaa gagcactcaa gctcttcata 1740
 acaccaacct gaccagagaa gtcaggcccc tctctgcctc gcagccttcg ctgccccatg 1800
 aggtctccac tatgatctcc agaccgcatc ccactgtggg cgagtccttg gcctccatcc 1860
 ctcaaccctg ggcaacagtc cacagcactg gccttcaggc agggagcagg agcaccgtgc 1920
 cacagcgtgt caccttggtc agacagatgt cctcgggagc tattttcccc aaccgaggag 1980
 tgccctccagc acccccacca ccagcagctg tgcagagaga gtctccctca gtcttaataa 2040
 aagaccaga tgcagaaaaa ccacgttttg cttcgaattt atgattcttg ctgattgtca 2100
 aagcagaaaa gaaatactct aataaacaga atattctcag atattatattt attctatctc 2160
 atgatagagc cctatagcct actctaaaaa gatatttttag aagctctggc gtacatgcaa 2220
 atgtaaaaac atatatacat atattattaa atatatatat atatctaaat gcccaagaga 2280
 agttcaaaag acttgataaa ctttcagtgat tatgtcttcc tttcttttaa accattaaag 2340
 gatttaacac attgttgtaa gatcattgat ttctaacctt ttacttaatt cctttgttat 2400
 atgtgtttct cctttttatg aagagttctt gaagtcattg gaaacaaaac tctgatttag 2460
 aaataaaaagg caactccaat tagtttcagc atagcaccaa tcaaagcttt ctttcattaa 2520
 ctgtgcctct gcatctaggt tgtaatttat gtgggattca ataaagaaat cccagtttat 2580
 agctctaaat tgtattttgg tgctttaaat tttgagttat gtgaaggaaac acactacacg 2640
 ctgagccacc ataggagact aacattgccca ctgttaaggc ttcctctaac ctcaaacatg 2700
 ttcgtcaatt ttgtgaggaa aggtgaggag atatttgtct tcatgtgtta ttggactttt 2760
 accaagattc agtcaatgtt agctgtaaat aacttttcca acctgaataa aagtaactat 2820
 tctgtgttgt ataaaggtaa aagtcactgt ttaagaattt agttttattg cttcacttca 2880
 aaagttagag ttttaaaatt tcacaaaaca taataattgt gacaactgtt caaatgtaat 2940
 gcaattgctt gagacctaca atatcattta aacctgcaat attttatgca aaaattgtat 3000
 gcttgaacct acaaatgtct tgtattacac caaaaatcat tacttttatt ccttcttgac 3060
 ataatcaagc atctgaacct agtcctggca tgcttttggg ggcaaaaaaa aa 3112

<210> 3

<211> 2606

<212> DNA

<213> Bos taurus

<400> 3

cgggagcccc gagcgcagcc actgagggca gcggcgccgg cgggagcgag gcgcgcagcg 60
 agaagcggcg gcgaggaatc ggccggggggc ttcgaggacg ccgagggggc ccggcgccag 120
 tacggcttca tgcagcggca gttcacctcc atgctgcagc ccgggggtcaa caaattctcc 180
 ctccgcagt tccggagcca gaaggcggtg gagaaggagc agggaaagggt taaaactgca 240
 ggcttctgga ttatccacc ttacagtgat ttcaggtttt attgggattt aataatgctt 300
 ataattgatg ttggaaatct ggtcatcata ccagttggaa tcacattctt tacagaacag 360

acaacaacac catggattat tttcaatgtg gcttcagata cagttttcct tttggacttg 420
 atcatgaatt tcaggactgg gactgtcaat gaagacagtt ctgaaatcat cctggaccct 480
 aaagtgatca agatgaatta tttaaaaagc tgggttgggg ttgacttcat ctcatacaatc 540
 ccagtggatt atatctttct cattgtagaa aaaggaatgg attcgggaagt ttacaagaca 600
 gccagggcac ttcgcattgt gaggtttaca aaaattctca gtctcttgcg tttattacga 660
 ctttcaagggt taattagata catacatcag tgggaagaga ttttccacat gacatatgat 720
 cttgccagtgt ctgtgggtgag aatttttaac ctcatggca tgatgctgct cctgtgccac 780
 tgggatggct gtcttcagtt cctggtagca ctgctgcagg acttcccacc agattgctgg 840
 gtgtctctaa atgagatgggt taatgattct tggggaaagc agtattccta cgcgctcttc 900
 aaagcgatga gtcatatgct gtgcattggc tacggagccc aagccccgt gagcatgtct 960
 gacctgtgga tcaccatgct gagcatgac gtcggggcca cctgctacgc catgtttgtt 1020
 ggccacgcca cggtctaat tcagtctttg gattcctcaa ggcggaata tcaagagaag 1080
 tataagcaag tggacaata catgtcattc cataagttac cagctgatat gcgtcagaag 1140
 atacatgatt attatgaaca cagataccaa ggcaaaatct ttgatgagga aaatattctc 1200
 aatgaactca atgactctct gagagaggag atagtcaact tcaactgccg aaaactagt 1260
 gctacaatgc ctctttttgc taatgcggat cctaatttcg tgaccgccat gctgagcaag 1320
 ttgagatttg aggtgtttca acctggagat tatatcatalc gagaaggagc tgtggctaaa 1380
 aaaatgtatt tcattcaaca tgggtgttgc ggtgtcatca caaaatccag taaagaaatg 1440
 aagctgacag atggctcata ctttggagag atttgcttgc tgaccaaggg acggcgact 1500
 gccagtgttc gagctgatac atattgtcgt ctttactcac tttctgtgga caatttcaat 1560
 gaggtcctgg aggaatatcc aatgatgaga agagcctttg agacggttgc cattgaccga 1620
 ttagatagga tagggaagaa aaattcaatt ctcttgcaaa agttccagaa ggatctgaac 1680
 acgggtgttt tcaacaatca ggagaacgag atcctgaagc agattgtgaa acacgacag 1740
 gaaatgggtgc aggcaatccc tcccctcaat taccctcaaa tgacagccct gaattccacc 1800
 tcttcaacta ctaccccgac ctctcgctg aggacacagt caccgccagt gtacacagcc 1860
 accagtctgt ctcatagcaa cctgcactcc cccagcccca gcaccagac ccccagccg 1920
 tcagccatcc tctcgccctg ctctacacc accgctgtct gcagccctcc tgtacagagc 1980
 ccgctagcca ctcgaaactt cactatgcc tccccacgg cttcccagtt gtccctcatt 2040
 cagcagcagc aggttcagca gccaccgcag cccagcagc caccccaacc tccacagacc 2100
 cccggcagct ccacaccgaa aaacgaagt cacaagagca cgcaggcgct tcacaacacc 2160
 agcctgaccc gagaagtcat gcccctctcg gcctcgagc cctcgctgcc ccacgaggtc 2220
 tccaccctga tctccagacc gcatcccact gtgggagag ccctggcctc catccctcaa 2280
 cccgtgacca cgggtccagg ctcgggcctg caggcagggg gcaggggac cgtccccag 2340
 cgagtcaccc tgttccgaca gatgtcatcg ggagccatcc ccccaatcg aggagtcccc 2400
 ccggccccc ctccaccagc agccgctcat ccgagggagg cgcctcagt cttaactaca 2460
 gactcagagg cagaaaagcc acgatttgc tcaaatttat gatcctgctg attgtaaagc 2520
 agaaagaaat actctaact aactgaggac gcttctcaga tttgatttta ttctatctcc 2580
 tgatagatcc tctagcctac tatgaa 2606

<210> 4

<211> 2986

<212> DNA

<213> Strongylocentrotus purpuratus

<400> 4

cgggagaata gtgcaccaag ggatgcccgt gaaatattaa ttaaacgttt ttaagaacat 60
 catcaaacc gggcccatc atgaaggaat aacaaggcct tcgaaaagta tgggaaactg 120
 gtcggcagga catcagcatt attaatctca ggaaactcat tatggataac aaggaaacta 180
 acggagagct agagcagctc gatgaggccg atccgtccgg tcaaaacctt gatgatgggg 240
 aaaccgatag caaacaagaa gagaatctca tcaacgttag cccgccaaaa acaccgccag 300
 gtctctctcc tctctctaaag aatggaggaa ggggtcagaa accgccaaa atcccaatat 360
 gtcatacaaaa tggaaagctc cccaaggaag ttgaatggac agaagacaga ggcaagaca 420
 gaaaggatag tctcactctt caatcaaagc tagatcacgg ggcatacacg gatgagaaac 480
 aggatcttct aacatatctt gaccgtcacg gcatcaacag tccagtcaag ctaacaccag 540
 atgaaactgg agggagcagt gctttggata ttcttgggat tattgaagag agggacactg 600
 gtgcactagg ctctgatccc tcatccacta tgcaggccat ggctaaacct gtaggctttc 660
 tgcagaggca gctatggact gtctcccaac cttcagacaa tagactctcc atgaaacttt 720
 tcggaagcaa gaaaggggta caaaaggaag aatatcggtc gaggaaggcg ggggttctta 780
 tcattcatcc atgtagtcatt ttcagatttt actgggatct actgatgctg tgctgatca 840
 tggcaaacgt catctccta ccgctcgtca ttactttctt ccacaacaag gacatgagta 900

cggggttggt catctttaat tgettctcag ataccttctt cattctcgat ctcatctgca 960
 acttttcggac cggcatcatg aatccgaagt cggccgaaca ggtgatcctc aacccccgtc 1020
 aaatcgcccta tcattatctc cgttcatggt tcacatcoga tctcgtgtct tccatcccca 1080
 tggactacat ctctctctc gctggcgcc agaaccgtca ctctctcgag gtgtcccgag 1140
 cctcaagat actgcgctt gccaaagtcc tcagtcttct tcgactcctg cgtctgtcca 1200
 ggctcatgog gttcgtcagt caatgggaac aggccttcaa cgtagccaat gccgtcatcc 1260
 ggatctgtaa tctagtgtgt atgatgttc tgattggcca ttggaatggc tgccttcaat 1320
 atctcgtgcc catgctgcaa gaatacccg accaatcatg ggtcgccatt aatggccttg 1380
 agcacgctca ttggtgggag cagtatacat gggcactctt caaagccctt tcgcacatgc 1440
 tctgtatcgg gtacggcaag tccccctc aaagcatcac cgatgtctgg ctaacgattg 1500
 tcagtatggt gtccggtgog acctgcttcg cctgttcat cggacacgct accaatctca 1560
 tccagtccat ggactcctcc agcaggcaat accgtgagaa gttgaaacaa gttgaagagt 1620
 acatgcagta tcgcaagcta ccgtcccacc tacgaaacaa gatcctcgat tactacgagt 1680
 accgataccg aggaagatg tttgatgaga ggcatactt tcgagaagtg tcggagagta 1740
 tacgacagga tgtcgcaaac tacaattgtc gcgacctggt cgcacccgtc cctttcttcg 1800
 tcggtgccga ctcaaacttc gtcacccgtg tggtagcgt gctcgaattc gaggtcttcc 1860
 aaccgctga ctatgttata caggaaggta ctttcggtga tcgcatgttc ttcacccagc 1920
 agggcatcgt cgacatcatc atgtccgacg gcgtcatcgc cactgcactc agtgacggct 1980
 catatttttg cgaaatctgc ctgcttaccg gtgagcgccg cgtggcatcg gtgaagtgcg 2040
 agacctactg cagctcttc tcgctctccg tccagcattt caaccaagtg ctcgacgagt 2100
 ttcccgccat gaggaacacg atggaagaga tagccgttcg tcgtctgacc cgaatcggga 2160
 agaatcgag caagctgaaa tccgcctag agagcccgac gatcaggggac actgccccctc 2220
 tctttccgat cccacctgat acacgtctt tcgtcaccga catcgaaaag aaccggttct 2280
 ttggcgacga cagcgacgat gtacacatca ggacccgagt cgacgtcgag cgtggttcgc 2340
 atgaaaacgt catcgccatc atggatggga gtttatccga cctcaggatg gaaaacgaaa 2400
 tccaagcccg taaatcgtct agcggaaaac ggaggaaatt ccagcaacaa acaaccgaac 2460
 tatgacgact tgaaacaaac aatgatggac gcttacaatt tccagtgatt caatacttac 2520
 gcaatgcaga cattagcttt tgtacctgat tgtttagaat gtattgaatt tgtagatcag 2580
 tccggcaaat aagaaagcat aatttggaa tttctttcatt gaggaagtac tgaaaacaat 2640
 gtgatagcag ccggtagaaa tttctgtgcc attatcgagg ctatatcttt cgcgcttttct 2700
 tacgaagtaa atgaaaggat caattaaaatt attgttcttt gtctcgtgcg ctttgtatct 2760
 gatgccgaaa agaatgaaa cgtgattaga acagtaatcg attgaactac agaagtcttt 2820
 tcaaaatggt gaatgtatga aggaggaggg ggaaggtttg atatatgcaa agaaatggag 2880
 aaatattttt gtaaatattat ctagaatggt actattgatg ctggaaagggt gttgaagttg 2940
 tccaatattg tgtcaaatca ccaactattt gacatttgtc tttttc 2986

<210> 5

<211> 3185

<212> DNA

<213> *Drosophila melanogaster*

<400> 5

cggaaatttcc tcgctgaagg gcaaggggca gagtcagagt caggggagaga gcggcagacg 60
 ctgcccggcc atcgcggtgc ggtgaggagc gagagtggaa gcgggagcag ccacaccatt 120
 ccggcgacgg gcaagagtc gccggtgccg cactcgtctg cggccaagat cagcagctcg 180
 gcaagcggca gcaagaactg caatttgtc agcgccagca gcaactcatg ccacaagctg 240
 aacgcccacg cccaaggatc ggagcaggat cgggatcttg gatcgggatc aggatcagga 300
 ccacccggac acagtcacta cgcggccgcc tcgccccaaa gctcggtcag cagcaacggt 360
 catctgaaca agtactgct cagcgacctc acgogccgca acgogagttc aatcgccagc 420
 tgagcgcgcc cagcgactac acgcaccact cctccagcaa cggatcgag caggaggggt 480
 cctcggaggc caacgagggc caccgaaccg tcggcgagtc caccatcacc gtagccagt 540
 ccggcgatc gtatccgcat ccgtactcct atccgtatca ttacggcacc accgctcctc 600
 ggccacagcg ccggccaatc tcaaggcgtc gctgcagctg cacagctttg ggagccacca 660
 tccgtgtcct tatccggcaa ggcccaacgt cagctcgtgc accaacagct tcaaccggcg 720
 ccacattcgc cggcacaagg gcaagctcgg cgatcgactg ctgagcgggg atagttagga 780
 atcgggtgcg tgctcctatt gctcgggtgt gaatgcgaac gacaacgacc tgcgcatttc 840
 gttcgagaac acctgcaccg attcgtggt aaccgcttct gatgatgaag cctgtcta 900
 atgcgaccaa ggaaccgaaa tggtagactt tgatgacgtg tcgttgtagc gactccgaa 960
 agaggagccc atgccaaca taccgatcgt gtcggaaaaa gtctctgca atttcctaaa 1020
 aagtcaattg caatcatggt tccagccgac ggacaaccga ctggccatga aactgtttg 1080

cagccgaaa ggcgtggtca aggagcgcac acgtcagaaa acttccgggc actgggtcat 1140
 acaccggtgc agttcattca ggttttactg ggacctttgc atgcttttat tattagtagc 1200
 aaatcttatt atcctgccag tcgcaatatc attcttcaac gatgatctga gcacacgatg 1260
 gattgccttc aactgcctaa gtgatactat ttttttaata gatattgtag tcaatttttag 1320
 aacaggaatt atgcaacaag acaacgctga acaagtaata ttggatccaa agcttatagc 1380
 taaacactat ttaagaactt ggttttttct cgatttgatt tcgtcgatac cgctagatta 1440
 tatattttta attttcaatc aaattatgaa attgcaggat ttctctgatt cttttcaaat 1500
 attgcatgcc ggacgcgccc tgcgatcct gcgcctggcc aagctgttat ccttgggtgcg 1560
 actgctccgc ctttcccgcc tcgtccgcta cgtttcccaa tgggaggagg tctatttctc 1620
 caatatggcc tcggtcttca tgaggatctt caatttaatt tgcattgatgc tctgatcgg 1680
 ccattggagc ggttgcttgc agttcttagt gccaatgttg caggggtttc catccaactc 1740
 ctgggtctcc atcaacgagt tgcaggaatc gtactggctg gagcagtatt cgtgggcatt 1800
 gttcaaggcc atgtcgaca tgcctctgat aggtacggc agattccgc cacaatcact 1860
 gacagacatg tggctgacga tgcctatgat gatatccggg gccacctgtt acgcattgtt 1920
 cctcggtcac gcgaccaatc tcatccagag cttggactcc agccggcgcc agtatcgca 1980
 gaaggtcaaa caggtggagg agtacatggc ctaccgcaag ctgccacgcg acatcgcgca 2040
 gcgcacacg gaatatttcg agcatcggta ccagggtaaa ttcttcgatg aagagttgat 2100
 acttggcgag ttgagcga aaactgcgca ggatgtcatc aactacaact gcagatccct 2160
 cgtggcgta gtgccttttt ttgctaattgc cgattcgaat ttctgttccg acgtagttag 2220
 caaactgaaa tacgaagttt tccaaccagg tgatattatc ataaaggagg gtacgatcgg 2280
 tactaagatg tacttcatac aggaggcggt ggtggacatt gtcattggca acggcgaggt 2340
 tgccacctca ctttcggatg ggtcttattt cggtgagatc tgcctgctga ccaatgcgcg 2400
 tcgtgtggcc agcgtgcgag ccgaaaccta ttgcagtcta ttctcgttga gcgtggatca 2460
 tttcaattgc gttctggatc agtatccgct gatgcgcaag accatggaga ctgtggccgc 2520
 cgagcgggta aacaagatcg gcaagaatcc aaacataatg catcagaagg acgagcagct 2580
 gagcaatccg gagtcgaaca cgattacggc tgtggttaat gcactggctg ccgaggcgga 2640
 tgactgcaaa gatgatgaca tggatctcag ggagaattta ctgcatgggt cagagtcgag 2700
 cattgctgag ccggtgcaga cgatacgtga gggctctccc aggccacgga gcggggagtt 2760
 ccgggccttg ttcgagggta acactccatg acactgagga gcagtgcaca gcggtgccct 2820
 cgggcaccgg gcaaccatct gaagcagcag ttcgctggac actcactcac caagtccac 2880
 atccatactc cacacaggac taccactcac acacacacac acactgcgta tataataatt 2940
 tagtaaaagg aaccccaaga cgcgataaga gtacactaaa aaaagaatca atttatggta 3000
 gacactctat atatgcaatt gcgatttagt agaaaacgta ttaaaaaacta aaaacccaaa 3060
 aaaagaagat aaaaacaatt acacaaaaaa tgcctcaat aattattcat aatttcagct 3120
 ccgctaactg tgatgacttt aatataagaa tcgaaaaaaa aattaacaaa caaacaaaaa 3180
 aaaag 3185

<210> 6

<211> 2922

<212> DNA

<213> Bos taurus

<400> 6

cgggagccc gagcgcagcc actgagggca gcggcgccgg cgggagcgag gcgcgcagcg 60
 agaagcggcg gcgaggaatc ggccgggggc ttcgaggacg ccgagggggc ccggcgccag 120
 tacggcttca tgcagcgcca gttcacctcc atgctgcagc ccggggtcaa caaattctcc 180
 ctccgcatgt tcgggagcca gaaggcggtg gagaaggagc aggaagggt taaaactgca 240
 ggcttctgga ttatccaccc ttacagtgat ttcaggtttt attgggattt aataatgctt 300
 ataattgatg ttggaaatct ggtcatcata ccagttggaa tcacattctt tacagaacag 360
 acaacaacac catggattat tttcaatgtg gcttcagata cagtttttct tttggacttg 420
 atcatgaatt tcaggactgg gactgtcaat gaagacagtt ctgaaatcat cctggaccct 480
 aaagtgatca agatgaatta tttaaaaagc tgggttgtgg ttgacttcat ctcatcaatc 540
 ccagtggtat atatctttct cattgttaga aaaggaatgg attcggaagt ttacaagaca 600
 gccagggcac ttcgatttgt gaggtttaca aaaattctca gtctcttgcg tttattacga 660
 ctttcaagggt taattagata catacatcag tgggaagaga ttttccacat gacatatgat 720
 cttgccagtg ctgtgggtgag aatttttaac ctcatggca tgatgctgct cctgtgccac 780
 tgggatggct gtcttcagtt cctggtacca ctgctgcagg acttccacc agattgctgg 840
 gtgtctctaa atgagatggt taatgattct tggggaaagc agtattccta cgcgctcttc 900
 aaagcgatga gtcattatgct gtgcattggc tacggagccc aagccccct gagcatgtct 960
 gacctgtgga tcaccatgct gagcatgatc gtcggggcca cctgctacgc catgtttgtt 1020

09640582.081700

ggccacgcca cggctctaata tcagtctttg gattcctcaa ggccggcaata tcaagagaag 1080
 tataagcaag tggacaata catgtcattc cataagttac cagctgatat gcgtcagaag 1140
 atacatgatt attatgaaca cagataccaa ggcaaaatct ttgatgagga aaatattctc 1200
 aatgaactca atgacccctc gagagaggag atagtcaact tcaactgccg aaaactagt 1260
 gctacaatgc ctctttttgc taatgcggat cctaatttcg tgaccgccat gctgagcaag 1320
 ttgagatttg aggtgtttca acctggagat tatatcatac gagaaggagg ctgtggtaaa 1380
 aaaatgtatt tcattcaaca tgggtgttgc ggtgtcatca caaaatccag taaagaaatg 1440
 aagctgacag atggctcata ctttggagag atttgcctgc tgaccaaggg acggcgact 1500
 gccagtgttc gagctgatac atattgtcgt ctttactcac tttctgtgga caatttcaat 1560
 gaggtcctgg aggaatatcc aatgatgaga agagccttg agacggttc cattgaccga 1620
 ttagatagga taggtactgt ttattttctt ctttacttac aattcacttt taactagt 1680
 gttgagtata ttttgcagt cataagtcac aaatgctagt ttacagattg cttattaact 1740
 agcatagaaa cagcaattag ctgtagccat atttctagaa gatctgaggc actaacttct 1800
 cgtctaagta ttctaggttt gtttattcat ctctgttttt actagcttca cagtctgatt 1860
 tcctcagtga taccaaaagg taaaaccaat gattacaaat tctagatggc attaaaatag 1920
 wwctnaaaaa tacaatagta tgagtctaca ttacaaacta tttttatna caagtttttt 1980
 ttttaanntt aagggtcaac attacattta ttcttatatt aagaattgaa aagaattgtg 2040
 cattttactt gtcacagtag aaacgttaat gtttgaata crrrctcaag cagaaaaagc 2100
 cttaatagaa ctgcccacat agatgcttta ttttgcaaac atcaacttat tttaaaatct 2160
 ttctgtctct caaattaaaa tattgatata taaggcctta ctagtattac tagtttaaac 2220
 gtctgaataa ttgccatgta aaaattagat cagattggct tgctgttaac ttcccaagat 2280
 atgctggaac attctgatgt cagaagggtg tatgcattca ttttccacac ccaaattctc 2340
 ctccccgacc agacccttct ctgctccctt tcccagctta actctactag ccttcatagt 2400
 tcaattttaa catcatttcc ctgtagaaac cnatngacct tccactcctc cttaatrta 2460
 tgagcaccct ggatattgtt tncataccc ctgggatgtt cctccatcac agtacagntt 2520
 ttattattta aattgctcta gagatnctaa gctttatgaa tnaagagatc angctcaatt 2580
 cactattaca ttcacagtag cwwgtacaca atgaatattg ttgaagagag ttagggaggg 2640
 atgaaggaat caatgaactc aaaggagatg ggggtgggat cactgaaaag taaacaaaga 2700
 ggtacttcaa ctgcttcatt cttattaaag gtaaggactt ttgattgatg ttacanttat 2760
 gtagctttt cttctgcact ttancatctt tcttttctc tatattagta ggacagaaga 2820
 ctgcataagg atctaggggt tgggttaggac aagtaaagggt agtatttggg cattaccatt 2880
 atggacacaa caaggcttcc aggtggataa caataataac gg 2922

<210> 7

<211> 1820

<212> DNA

<213> Bos taurus

<400> 7

cggcgacga ggccggcagc gaggaggcgg gcccgccggg ggagtcgcgc ggcagccagg 60
 ccagcttcat gcagcgccag ttccggcgcg tctgcagcc gggcgctcaac aagttctcgt 120
 tgccgatgtt cggcagtcag aaggccgtgg agcgcgacga ggagcgctt aagtcagcgg 180
 gggcctggat catccaccct tacagcgact tcaggttcta ctgggacttc accatgctgc 240
 tcttcatggt gggaaacctc atcatcatcc ccgtgggcat cacttcttc aaggacgaga 300
 ccacggcccc atggattgtg ttcaatgttg tctcgacac attcttctc atggacctg 360
 tgctgaactt ccgcacgggc attgtgatcg aggacaacac ggagatcatc ctggaccccc 420
 agaagatcaa gaagaagtac ctgcgcacgt ggttcgtggt ggacttcgta tctccatcc 480
 ccgtggtaag ctacatcttc ctcatcgtgg agaaaggcat cgactctgag gtctacaaga 540
 cggcccgcg cctgcccgatc gagccgttca ccaagatcct cagcctgctg cgctgctcc 600
 gcttgtcgcg cctcatccgc tacatccatc agtgggagga gatcttcac atgacctacg 660
 acctggcgag cgcctcatg cgcctctgca acctcatcag catgatgctg ctctctgccc 720
 actgggatgg ctgctgcag ttctgtgtgc ccatgcttca ggacttccca cgcaactgct 780
 ggggtctccat caacggcatg gtgaacct catggagcga gctctactcc ttccgctgt 840
 tcaaggccat gagccacatg ctgtgcacg ggtacgggcg gcaggcgcca gaaagcatga 900
 cggacatctg gctgacctg ctgagcatga tctgggtgc cactgctac gccatgttca 960
 ttggccacgc caccgcccctc atccagtcg tggactcctc aaggcgccag taccaggaga 1020
 agtacaagca agtggagcag tacatgtcct tccacaagct gccagccgac ttccgcccaga 1080
 agatccacga ctactacgag caccgctacc agggcaagat gttcgacgag gacagcatcc 1140
 tcggcgagct caaggcgggc ctgcccggag agatcgtcaa cttcaactgc cggaagctgg 1200
 tggcctccat gccactgttc gccaatgctg accccaactt cgtcacgggc catctgacca 1260

```

agctcaagtt tgaggtcttc cagccaggcg actacatcat ccgtgagggc accattggca 1320
agaagatgta cttcatccaa cacggcgtgg tcagtgtgct taccttgggc aacaaggaga 1380
tgaagttgtc tgatggctcc tactttgggg agatctgctt gctgacgcgg ggccggcgca 1440
cggcgagcgt ccgggcccgc acctactgcc gcctctactc gctgagtgtg gacaacttca 1500
atgaggtgct ggaggagtac cccatgatga ggccgggcctt tgagacagtc gccattgacc 1560
gcctggatcg cattggcaag aagaactcga tcctgctaca caaggtgcag cacgacctca 1620
actctggcgt gtttaacaac caggagaacg ccatcatcca ggagattgtc aagtatgacc 1680
gcgagatggg gcagcaggct gagctggggc agcgtgtcgg cctcttcccg ccaccaccgc 1740
cacctccaca gggcacctca gccattgcca cgctgcagca gccgtggcca tgagcttctg 1800
tccacaagtc gcacgcccc 1820

```

```

<210> 8
<211> 101
<212> DNA
<213> Rattus rattus

```

```

<400> 8
ctacatcatc cgagagggga ccatcgggaa gaagatgtac ttcattccagc acgggggtgg 60
gagcgtgcta accaggggca acaaggagga taagctgtca n 101

```

```

<210> 9
<211> 558
<212> DNA
<213> Rattus rattus

```

```

<400> 9
tctgggtggg cgtgagggct ccgtggggcag gaagatgtac ttcattccagc atggcgtgct 60
cagtgtgttg gcacggggcg ctccgggacac tcgcctcact gacggatcct actttgggga 120
gatctgcctg ctgactcgag gtcggagaac agccagtgtg agggctgaca cctactgtcg 180
cctctactca ctcagcgtgg accacttcaa tgcagtgtct gaggagctcc cgatgatgcg 240
cagggccttt gagactgtgg ccatggaccg gcttcggcgc atcggtgagg cctgtctgcc 300
ctgtctgtct tgggcccctg ctgagcctca tctcattttc atagcaagga acctaccct 360
agtgtttctt ctccacaccc caacctaccc agtaccagca ggctattagc tctgtttctc 420
gctagtctta cccctagaaa gaaatagcca tggagctgtc tccccaaacc ctcatccct 480
gtgtcctctc gggtagcagt acttaacctc accgtttttg ataccacctt ccagtttctg 540
ttgccaagca ttctctcc 558

```

```

<210> 10
<211> 2886
<212> DNA
<213> Homo sapiens

```

```

<400> 10
gaattcgcgg ccgcgtcgac ggccagcttc atgcagcgcc agttcggcgc gtcctgcag 60
ccgggcgtca acaagttctc gctgcggatg ttccggcagcc agaaggccgt ggagcgcgag 120
caggagcgcg tcaagtcggc gggggcctgg atcatccacc cgtacagcga cttcagggtc 180
tactgggact tcaccatgct gctgttcatg gtgggaaacc tcatcatcat ccagtgggc 240
atcaccttct tcaaggatga gacctgccc ccgtggatcg tgttcaacgt ggtctcggac 300
accttcttcc tcatggacct ggtgttgaa tcocgcacog gcattgtgat cgaggacaac 360
acggagatca tcctggaccc cgagaagatc aagaagaagt atctgcgcac gtgggtcgtg 420
gtggacttcg tgtcctccat ccccgctggc tacatcttcc ttattgtgga gaagggcatt 480
gactccgagg tctacaagac ggcacgcgcc ctgcgcacog tgcgcttcac caagatcctc 540
agcctcctgc ggctgctgcg cctctcacgc ctgatccgct acatccatca gtgggaggag 600
atcttccaca tgacctatga cctggccagc gcgggtgatg ggatctgcaa tctcatcagc 660
atgatgctgc tgctctgcca ctgggacggc tgccctgcagt tcctgggtgcc tatgctgcag 720
gacttccgcg gcaactgctg ggtgtccatc aatggcatgg tgaaccactc gtggagtga 780
ctgtactcct tcgcactctt caaggccatg agccacatgc tgtgcacog gtacggccgg 840
caggcgcccg agagcatgac ggacatctgg ctgacctgac tcagcatgat tgtgggtgcc 900
acctcgtacg ccatgttcat cggccacgcc actgcctca tccagtcgct ggactcctcg 960
cggcgccagt accaggagaa gtacaagcag gtggaacagt acatgtcctt ccacaagctg 1020

```

002180 28504950

ccagctgact tccgccagaa gatccacgac tactatgagc accgttacca gggcaagatg 1080
 tttgacgagg acagcatcct gggcgagctc aacgggcccnn tgcggnagga gatcgtcaac 1140
 ttcaactgcc ggaagctggt ggcctccatg ccgctgttcg ccaacgccga ccccaacttc 1200
 gtcacggcca tgctgaccaa gctcaagttc gaggtcttcc agccgggtga ctacatcatc 1260
 cgccaaggca ccatcgggaa gaagatgtac ttcattccagc acggcgtggt cagcgtgctc 1320
 actaagggca acaaggagat gaagctgtcc gatggctcct acttcgggga gatctgcctg 1380
 ctaccccggg gccgcgcac ggcagcgtgc gngctgacac ctactgccgc ctctatttcg 1440
 tgagcgtgga caacttcaac gagntgctgg aggagtacc ccatgatgcgg cgcgccttcg 1500
 agacggtggc catcgaccgc ctggaccgca tcggtgagcg ggcggggggc gtggccgggg 1560
 cgggtgccct ggcggggggag gggcgtggcc aaggcatcag gagagtggct tggacagtgg 1620
 cagggggaag ggcgtggctg tggcatcagg ggcacgggtg ggcagagac gtggccaagg 1680
 catncaggag tgtggccatg gcagcagggg cgtggctggg gcaggggcag cggctggccg 1740
 ctctaggac ccttttgggt cttagaggctg attttctgac ctattgtcct acttcagcca 1800
 gaggcagcct gtttcccaag ggaggggaatg cacaggggtg ttgcggttgt gccgaatgct 1860
 cggtgagcac ctgctgtgtg ctgggggtgc aggggacaga cccggggggc cactcagact 1920
 cccagggagg cttatggact ggtgatgaaa tcacacacga ctgggctgtg tgccagcagg 1980
 gcaggtgggg ccggtgggct tccctgagtt gggaatgcag agtggagacc agggtaaggg 2040
 atgccatgtg gaaacgggga ggaagatgtg ttcgtggagt ggacacagca catcccaagg 2100
 ccctgaggtg gaaaagaggc cttagagtcca gagagccagg gaggcctgga ggaggttggg 2160
 gaagaagggg agggcagaca cacagggccc agtgggcggc agggagagtt tagactaaat 2220
 caggagcatc agggagccat ggaggggtct aggtgggcgg aggacctggt cagattgtat 2280
 ccgccaaggc gggccgtgtc caggagggag acggtgacct ggctctcag gggggcagtc 2340
 tctggggcag ggagggncag agccctgatg actggatgta ggccgagag agatggcggc 2400
 tcatgctgct gttcgtggga atgggaatga agaccatggc tgaaacgcag gacaggtgcg 2460
 acggagtggg gtcagggagc tccctggtgt acagtaggaa gctctccaca acttgctcta 2520
 tacagtgagt atgcaaccgc ttctgagta tcaggtgctt aggttataac ttctgtatac 2580
 agcaggtgct cagcacaggc tgtgtacagg caggtgtttt cggtatgcct gtggcacact 2640
 ggaggcagtc attacataat cagcgtatac aggtgggtaca catgcatact tgggtgcacag 2700
 tgatacctgc tccatgtaca cagcaggcat taaatacctg tttactgcca ggcgcggtgn 2760
 ntcacgcctg tagtcccagc actttcggag gccaaaggtg gtggatcacg aggtcaggag 2820
 attgagacca tccctggctaa catggtgaaa ccccgctctc actaaaaaaaa aaaaaaaaaa 2880
 aaaaaa 2886

<210> 11

<211> 2029

<212> DNA

<213> Homo sapiens

<400> 11

gcgngccgcg tcgacgtggc ctccatgccca ctgttttgcca atgcgggaccc caacttcgtg 60
 acgtccatgc tgaccaagct gcgtttttgag gtcttccagc ctgggggacta catcatccgg 120
 gaaggcaacca ttggcaagaa gatgtacttc atccagcatg gcgtgggtcag cgtgctcacc 180
 aagggcaaca aggagaccaa gctggccgac ggctcctact ttggagagat ctgctgctg 240
 acccgggggc ggcgcacagc cagcgtgagg gccgacacct actgccgcct ctactcgtg 300
 agcgtggaca acttcaatga ggtgctggag gagtacccca tgatgcgaag ggccttcgag 360
 accgtggcgc tggaccgcct ggaccgcatt ggcaagaaga actccatcct cctccacaaa 420
 gtccagcacg acctcaactc cggcgtcttc aactaccagg agaatgagat catccagcag 480
 attgtgcagc atgaccggga gatggcccac tgcgcgcacc gcgtccaggc tgetgnctct 540
 gccaccccaa ccccccacgc cgtnatctgg acccgcgtga tccaggcacc actgcaggct 600
 gccgctgccca ccacttctgt ngccatagcg ctaccccacc acccytcgcn tgnctgytgc 660
 natnttnncg sectncccc anggatctnn gggctggnc aacctcggtgc cggnmagang 720
 ccaaggcacc tagnaacggct gnagtnctg atcccttctg cgtgggtccg cctcgccgcg 780
 cagcagcccg tcccagggtg acacaccgct ttcactctcc tccacatcc aacagctggc 840
 tggattctct gcccccgtg gactgagccc actcctgccc tcatccagct cctccccacc 900
 ccccgggggc tgtggctccc cctcggtccc cacaccatca gctgcgtagc cgcaccacc 960
 atagccgggt ttggccactt ccacaaggcg ctgggtggct cctgtctctc ctccgactct 1020
 cccctgctca ccccgctgca gccaggcgcc cgtccccgc aggtgcccga gccatctccc 1080
 gcgccacccg gggcccgggg aggcctggga ctcccgagc acttctgccc acccccaccc 1140
 tcatccagat ccccgctatc tagccccggg cagctgggccc agcctcccg ggagttgtcc 1200
 ctaggctctg ccactggccc actgagcacg ccagagacac cccacggca gcctgagccc 1260

```

ccgtcccttg tggcaggggc ctctgggggn ggnttccctt gtaggncttt actccccgag 1320
gaggtntcag cccccstggn ccacagccna gsccccnaa gaaccttccc gagtgtcccg 1380
ccccggncnt ctggctccca crgantonnn ctryyctg ccacctgcat ccagcccccc 1440
accacccag ntccccagc gccgggncac acccccgctc acccccggcc gcctcaccga 1500
ggacctcaag ctcatctccg cgtctcagcc agccctgcct caggacgggg cgcagactct 1560
ccgcagagcc tccccgcact cctcagggga gtccatggct gccttcccg ccttccccag 1620
ggctgggggt ggcagcgggg gcagtgggag cagcgggggc ctcggtcccc ctgggaggcc 1680
ctatgggtgc atccccggcc agcacgtcac tctgcctcgg aagacatcct caggttcttt 1740
gccaccccc ctgtctttgt ttggggcaag agccacctct tctggggggc cccctctgac 1800
tgctggaccc cagaggggaa ctggggccag gtgcgtcca aactgccgtc 1860
caatctatga gctgggccct tccttccctc ttctttcttc tttctctccc cttccttctt 1920
ccttcagggt taactgtgat taggagatat accaataaca gtaataatta tttaaaaaac 1980
cancasacac cagaaaaaca aaagacrrnc agaaagtcga cgcggccgc 2029

```

<210> 12

<211> 2984

<212> DNA

<213> Bos taurus

<400> 12

```

gggcaccagc cgcgccggag cccggagcgc agccactgag ggcagcggcg gggcggggag 60
cgaggcgcgc agcgagaagc ggcgcggggc ggaagcagaa gccgccgccg ccgccgccgc 120
cgccgcgacg ggcagccggg ctccggcgcc cccggatcgg gccctgccc cctccgcctc 180
gtgtccccgg cgcggggcgg ccggcgagtc tggagccgc gccgtcggc gccgcgtccc 240
ccgggcatgg aaggaggcgg caagcccaac tcctcgtcca acagccggga cgatggcaac 300
agcgtcttcc ccaccaaggc gccgcgacg ggcgcggggc cggccgcggc cgagaagcgc 360
ctgggcaccc cgcggggggg cggcgggacc ggcgcgaagg agcacggcaa ctcagtgtgc 420
ttcaagggtg acggcgggcg cggcgggcgc gaggaatcgg ccggggggctt cgaggacgcc 480
gaggggcccc ggcggcagta cggcttcatg cagcggcagt tcacctccat gctgcagccc 540
ggggtcaaca aattctccct ccgcatgttc gggagccaga aggcggtgga gaaggagcag 600
gaaagggtta aaactgcagg cttctggatt atccaccctt acagtgattt caggttttat 660
tgggatttaa taatgcttat aatgatggtt ggaaatctgg tcatcatacc agttggaatc 720
acattcttta cagaacagac aacaacacca tggattattt tcaatgtggc ttcagataca 780
gttttccttt tggacttgat catgaatttc aggactggga ctgtcaatga agacagttct 840
gaaatcatcc tggaccctaa agtggattat cgcatttgta ttgtagaaa aggaatggat 960
gacttcatct catcaatccc cagggcactt cgtatttgta ggtttacaaa aattctcagt 1020
tcggaagttt acaagacagc cagaatgact ttcaagggtta attagataca tacatcagtg ggaagagatt 1080
ctcttgctt tattacgact tgccagtgtc gtggtgagaa tttttaacct cattggcatg 1140
atgctgctcc tgtgccactg ggatggctgt cttcagttcc tgggtaccact gctgcaggac 1200
ttcccaccag attgctgggt gtctctaaat gagatggtta atgattcttg gggaaagcag 1260
tattcctacg cgctcttcaa agcgatgagt catatgctgt gcattggcta cggagcccaa 1320
gcccccgta gcatgtctga cctgtggatc accatgctga gcatgatcgt cggggccacc 1380
tgctacgcca tgtttgttgg ccacgccacg gctctaattc agtctttgga ttctcaagg 1440
cggcaatatc aagagaagta taagcaagtg gaacaataca tgtcattcca taagttacca 1500
gctgatatgc gtcagaagat acatgattat tatgaacaca gataccaagg caaaatcttt 1560
gatgaggaaa atattctcaa tgaactcaat gatcctctga gagaggagat agtcaacttc 1620
aactgccgaa aactagtggc tacaattgct ctttttgcta atgcgatcc taatttcgtg 1680
accgccatgc tgagcaagtt gagatttgag gtgtttcaac ctggagatta tatcatacga 1740
gaaggagctg tgggtaaaaa aatgtatttc attcaacatg gtgttgctgg tgtcatcaca 1800
aaatccagta aagaaatgaa gctgacagat ggctcatact ttggagagat ttgcttgctg 1860
accaagggac ggcgcactgc cagtgttoga gctgatacat attgtcgtct ttactcactt 1920
tctgtggaca atttcaatga ggtcctggag gaatatccaa tgatgagaag agcctttgag 1980
acggttgcca ttgaccgatt agataggata gggaagaaaa attcaattct cctgcaaaag 2040
ttccagaagg atctgaacac ggggtgtttt aacaatcagg agaacgagat cctgaagcag 2100
attgtgaaac acgacagggg aatgggtgcag gcaatccctc cctcaatta cctcaaagt 2160
acagccctga attccacctc ttcaactact accccgacct ctcgcctgag gacacagtca 2220
ccgccagtgt acacagccac cagtctgtct catagcaacc tgcactcccc cagccccagc 2280
accagacccc cccagccgtc agccatcctc tcgccctgct cctacaccac cgctgtctgc 2340
agccctcctg tacagagccc gctagccact cgaactttcc actatgcctc cccacggct 2400

```

```

tcccagttgt ccctcattca gcagcagcag gttcagcagc caccgcagcc ccagcagcca 2460
ccccaacctc cacagacccc cggcagctcc acaccgaaaa acgaagtgca caagagcacg 2520
caggcgcttc acaacaccag cctgaccoga gaagtcaggc ccctctcggc ctgcagccc 2580
tcgctgcccc acgaggtctc caccctgato tccagaccgc atcccactgt gggcgagtcc 2640
ctggcctcca tccctcaacc cgtgaccacg gtccacggct cgggcctgca ggcagggggc 2700
aggggcaccg tccccagcg agtcaccctg ttccgacaga tgtcatcggg agccatcccc 2760
cccaatcgag gagtcccccc ggccccccct ccaccagcag ccgctcatcc gagggaggcg 2820
ccctcagtct taactacaga ctgagaggca gaaaagccac gatttgcttc aaatttatga 2880
tccctgctgat tgtaaagcag aaagaaatac tctaactgtaa ctgaggacgc ttctcagatt 2940
tgatttttatt ctatctcctg atagatcctc tagcctacta tgaa 2984

```

```

<210> 13
<211> 794
<212> DNA
<213> Rattus rattus

```

```

<400> 13
tgccctgcagt tcctgggtgcc catgctgcaa gacttcccca gcgactgctg gtgtccatca 60
acaacatggt gaaccactcg tggagcgaac tctattcggt cgcgctcttc aaggccatga 120
gccacatgct ctgtattggc tacgggcggc aggctcccga gagcatgacg gacatctggc 180
tcaccatgct cagcatgata gtgggcgcca cctgctacgc tatgttcatt gggcacgcca 240
cggcgcttat ccagtccctg gactcgtcac ggccgagta ccaggagaag tacaagcaag 300
tggagcagta catgtccttc cacaaactgc cggctgactt ccgccagaag atccacgatt 360
actatgaaca ccggtaccag gggaagatgt ttgacgagga cagcatcctg ggggaactca 420
acggcccact gcgtgaggag attgtgaact tcaactgccg gaagctggtg gcttccatgc 480
cgttggtttgc caacgcagac cccaacttcg tcaccgccat gctgacaaag ctcaaatttg 540
aggtcttcca gcctggagac tacatcatcc gagaggggac catcggaag aagatgtact 600
tcatccagca cggggtggtg agcgtgctca ccaagggcaa caaggagatg aagctgtcag 660
atggctccta ttttggggag atctgcctgc tcacgagggg ccggcgca gccagtgtgc 720
gggctgacac ctactgtcgc ctctactcac tgagcgtgga caacttcaac gaggtgctgg 780
aggagtaccc catg 794

```

```

<210> 14
<211> 649
<212> DNA
<213> Rattus rattus

```

```

<400> 14
tccagcatgg gctgctcagt gtgttggcac ggggcgctcg ggacactcgc ctactgacg 60
gactctactt tggggagatc tgcttgctga ctgaggtcg gagaacagcc agtctaagg 120
ctgacaccta ctgtcgctc tactactca gcgtggacca cttcaatgca gtgcttgagg 180
agctcccgat gatgcgcagg gcttttgaga ctgtggccat ggaccggctt cggcgcatcg 240
gcaaaaagaa ttcgatattg cagcggaac gctctgagcc gactccaggc agcagcagtc 300
gtggcgtcat ggagcagcat ttggtacaac acgacagaga catggctcgt ggtattcggg 360
gtctggctcc gggcacagga gcccgcctca gtggaaagcc agttctgtgg gaaccactgg 420
tacacgcacc tcttcaggca gctgctgtga cctccaacgt ggccatagcc ttgactcatc 480
agcgaggccc tctgcccctc tcccctgatt ctccagccac cctcctggct cgatctgcta 540
gacgctcagc aggtcctcca gctcctccac tgggtgctgt tcgagcaggt cctctgctgg 600
cccggggacc ctgggcgtcc acttctcatc ttctgacca cgggcctc 649

```

```

<210> 15
<211> 4751
<212> DNA
<213> Homo sapiens

```

```

<400> 15
tcgacaaaaa tgccagggaa aggcgagccc agagcttggg gatggagaaa ttgggaagcc 60
acccccacc cttcaatctt aggatgggga attcgcaact gaagccggag cttcagactt 120
ggggcgcact cccagcttag cccaggaaag agatttaagg gcgcagcagt gtggatacct 180
ctcaccctcg cccgaaggt ctagcgaggg tctaactctg gcccttgcc aggcccgccc 240

```

```

ccccccctt  tccagccccc  ggcccggtgcg  ccgctgcccc  ttttaagaagc  ccaggtaggc  300
aggcccggt  gctggagccg  ctccataggg  aaccgcgag  ctgcggcggc  ttcataaata  360
ttccggggcg  cgggagcccg  agcgtgccc  gaggcgctt  cgggggaggc  ggccgctgat  420
gtaagcccg  cgggtcgctg  ggctccgctc  ggttgccggc  ggagcccccg  gacggggccg  480
acgggcccgg  gcagaggagg  cgaggcgagc  tcgcggtgg  ccagccacaa  agcccgggcg  540
gcgagacaga  cggacagcca  gccctcccgc  gggacgcacg  cccgggaccc  gcgcgggccg  600
tgcgctctgc  actccggagc  ggttccctga  gcgcgcggc  cgagagccct  ctccggcccg  660
cgccatttgt  tcccccgggg  ggccggggcg  ctggagcccg  gcggcgcgcc  gccctgaac  720
gccagggga  gggagggagg  caagaaggga  gcgcggggtc  cccgcgcca  gccggggccg  780
ggaggaggtg  tagcgcgcg  agcccgggg  ctggagccg  gactaggatc  ctcccccg  840
cgcgagccct  gcccaagcat  gggcgccgta  ggtgcccc  acgcggcg  caaaggacgc  900
gtccccacgg  gcggaactgac  cggcgggcg  acctggagcc  cgtccgcg  gccgcgctcc  960
tgcccccg  ccggtccgac  cccggccct  ggcgccatgg  acaagctgcc  gccgtccatg  1020
cgcaagcg  tctacagcct  cccgcagcag  gtgggggcca  aggcgtggat  catggacgag  1080
gaagaggagc  ccgaggagga  gggggccggg  ggccgccaag  accccagccg  caggagcatc  1140
cggctcggg  cactgccctc  gccctcccc  tcggcgccg  cgggtggcac  ggagtcccg  1200
agctcgggcc  tcggggcagc  ggacagcgaa  gggccggccc  gcggcgcg  caagtccagc  1260
acgaacggcg  actgcaggcg  ctcccgcg  agcctggcct  cgctgggcag  ccggggcg  1320
ggcacgggg  gcacggggag  cggcagcagt  cacggacacc  tgcatactc  cgcgaggag  1380
cggcggtca  tcgcccagg  cgacgcgtcc  cccggcgagg  acaggacgcc  cccaggcctg  1440
gcggccgagc  ccgagcgccc  cggcgccctc  gcgcagccc  cagcctcgcc  gccgcgccc  1500
cagcagccac  cgcagccggc  ctccgcctcc  tgccagcagc  cctcggtgga  caccgctatc  1560
aaagtggagg  gaggcgcg  tgccggcg  cagatccctc  cggaggccga  ggtgcgctg  1620
ggccaggcg  gcttcagca  gcgcagttc  ggggccatgc  tccaaccgg  ggtcaacaaa  1680
ttctccctaa  ggatgttcg  cagccagaaa  gccgtggagc  gcgaacagga  gaggtcaag  1740
tcggccggat  tttgattat  ccaccctac  agtgacttca  gattttactg  ggacctgacc  1800
atgctgctgc  tgatggtgg  aaacctgatt  atcattcctg  tgggcatcac  cttcttcaag  1860
gatgagaaca  ccacacctg  gattgtcttc  aatgtggtgt  cagacacatt  cttcctcatc  1920
gacttggtcc  tcaacttccg  cacagggatc  gtggtggagg  acaacacaga  gatcatcctg  1980
gaccgcagc  ggattaaaat  gaagtacctg  aaaagctggt  tcatggtaga  tttcatttcc  2040
tccatcccc  tggactacat  ctccctcatt  gtggagacac  gcatcgactc  ggaggtctac  2100
aagactgcc  gggccctgcg  cattgtccgc  ttcacgaaga  tcctcagcct  cttacgcctg  2160
ttacgcctct  ccgcctcat  tcgatataat  caccagtggg  aagagatctt  ccacatgacc  2220
tacgacctgg  ccagcgccgt  ggtgcgcac  gtgaacctca  tcggcatgat  gctcctgctc  2280
tgccactggg  accgctgct  gcagttcctg  gtacctatgc  tacaggactt  ccctgacgac  2340
tgctgggtgt  ccatcaacaa  catggtgaac  aactcctggg  ggaagcagta  ctccacgcy  2400
ctctcaagg  ccagagcca  catgctgtgc  atcggtacg  ggccggcagg  gcccggtggc  2460
atgtccgacg  tctggctcac  catgctcagc  atgatcgtg  gtgccacctg  ctacgccatg  2520
ttcattggcc  acgccactgc  cctcatccag  tccttgga  cctcccgcg  ccagtaccag  2580
gaaaagtaca  agcaggtgga  gcagtacatg  tcctttcaca  agctcccgcc  cgacaccgg  2640
cagcgcatcc  acgactacta  cgagcaccgc  taccagggca  agatgttcga  cgaggagagc  2700
atcctgggg  agctaagcga  gccctgcgg  gaggagatca  tcaactttaa  ctgtcggaag  2760
ctggtggcct  ccatgccact  gtttgccaat  gcggaccca  acttcgtgac  gtccatgctg  2820
accaagctgc  gtttcgaggt  ctccagcct  ggggactaca  tcatccggga  aggcaccatt  2880
ggcaagaaga  tgtacttcat  ccagcatggc  gtggtcagcg  tgctcaccaa  gggcaacaag  2940
gagaccaagc  tggccgacgg  ctccacttt  ggagagatct  gcctgctgac  ccggggcccg  3000
cgcacagcca  gcgtgagggc  cgacacctac  tgccgctct  actcgctgag  cgtggacaac  3060
ttcaatgagg  tgctggagga  gtaccccatg  atgcgaagg  ccttcgagac  cgtggcgctg  3120
gaccgctgg  accgatttg  caagaagaac  tccatcctcc  tccacaaagt  ccagcacgac  3180
ctcaactccg  cgtcttcaa  ctaccaggag  aatgagatca  tccagcagat  tgtgcagcat  3240
gaccgggaga  tggccactg  cgcgcaccgc  gtccaggctg  ctgcctctgc  caccacaacc  3300
cccacgccc  tcatctggac  cccgctgac  caggcaccac  tgaggctgc  cgctgccacc  3360
acttctgtgg  ccatagccct  caccaccac  cctcgctgc  ctgctgcat  cttccgccc  3420
ccccaggat  ctgggctggg  caacctcggt  gccgggcaga  cgccaaggca  cctgaaacgg  3480
ctgcagtccc  tgatcccttc  tgcgctgggc  tcgcctcgc  ccgcccagcag  cccgtcccag  3540
gtggacacac  catcttcatc  ctccctccac  atccaacagc  tggctggatt  ctctgcccc  3600
gctggactga  gccactcct  gccctcatcc  agctcctccc  cacccccgg  ggcctgtggc  3660
tccccctcg  ctccacacc  atcagctggc  gtaccgcca  ccacatagc  cgggtttggc  3720
cacttccaca  aggcgtggg  tggtccctg  tcctcctccg  actcctccc  gctcaccgc  3780
ctgcagccag  gcgcccgc  cccgcaggct  gccagccat  ctcccgcc  accgggggc  3840

```

cggggaggcc tgggactccc ggagcacttc ctgccacccc caccctcatc cagatccccg 3900
 tcatctagcc cggggcagct gggccagcct cccggggagt tgtccctagg tctggccact 3960
 ggcccactga gcacgccaga gacaccccca cggcagcctg agccgccgtc ccttgtggca 4020
 ggggcctctg ggggggcttc ccctgtaggc ttactcccc gaggaggtct cagccccct 4080
 ggccacagcc caggccccc aagaaccttc ccgagtgcc cgccccgggc ctctggctcc 4140
 cacggatcct tgcctctgcc acctgcatcc agccccccac caccaggt cccccagcgc 4200
 cggggcacac ccccgctcac ccccgccgc ctcacccagg acctcaagct catctccgcg 4260
 tctcagccag cctgcctca ggacggggcg cagactctcc gcagagcctc cccgcactcc 4320
 tcaggggagt ccattggctgc cttcccgctc ttcccagggt ctgggggtgg cagcgggggc 4380
 agtgggagca gcgggggcct cgggtcccc gggaggccct atggtgccat ccccggccag 4440
 cacgtcactc tgcctcgga gacatcctca gggtctttgc caccctct gtctttgttt 4500
 ggggcaagag ccacctcttc tggggggccc cctctgactg ctggacccca gagggaacct 4560
 ggggcccaggc ctgagccagt gcgtccaaa ctgccgtcca atctatgagc tgggcccctc 4620
 cttccctctt ctttctctt ttctctcct tcttcttcc ttcagggtta actgtgatta 4680
 ggagatatac caataacagt aataattatt taaaaaacca cacacaccag aaaaacaaaa 4740
 gacagcagaa a 4751

<210> 16

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerated Primer

<400> 16

ctgactgcag argtnttyca rccngnga 29

<210> 17

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerated Primer

<400> 17

atcggaattc nccraartan gancrtc 28

<210> 18

<211> 767

<212> PRT

<213> Strongylocentrotus purpuratus

<400> 18

Met Asp Asn Lys Glu Thr Asn Gly Glu Leu Glu Gln Ser Asp Glu Ala
 1 5 10 15

Asp Pro Ser Gly Gln Asn Leu Asp Asp Gly Glu Thr Asp Ser Lys Gln
 20 25 30

Glu Glu Asn Leu Ile Asn Val Ser Pro Pro Lys Thr Pro Pro Gly Pro
 35 40 45

Pro Pro Pro Leu Lys Asn Gly Gly Arg Gly Gln Lys Pro Pro Lys Ile
 50 55 60

Pro Ile Cys His Gln Asn Gly Lys Leu Pro Lys Glu Val Glu Trp Thr
 65 70 75 80

007120 22504360

65	70	75	80
Glu Asp Arg Gly Glu Asp Arg Lys Asp Ser Leu Thr Leu Gln Ser Lys	85	90	95
Leu Asp His Gly Ala Tyr Thr Asp Glu Lys Gln Asp Leu Leu Thr Tyr	100	105	110
Leu Asp Arg His Gly Ile Asn Ser Pro Val Lys Leu Thr Pro Asp Glu	115	120	125
Thr Gly Gly Ser Ser Ala Leu Asp Ile Leu Gly Ile Ile Glu Glu Arg	130	135	140
Asp Thr Gly Ala Leu Gly Ser Asp Pro Ser Ser Thr Met Gln Ala Met	145	150	155
Ala Lys Pro Val Gly Phe Leu Gln Arg Gln Leu Trp Thr Val Leu Gln	165	170	175
Pro Ser Asp Asn Arg Leu Ser Met Lys Leu Phe Gly Ser Lys Lys Gly	180	185	190
Leu Gln Lys Glu Lys Tyr Arg Leu Arg Lys Ala Gly Val Leu Ile Ile	195	200	205
His Pro Cys Ser His Phe Arg Phe Tyr Trp Asp Leu Leu Met Leu Cys	210	215	220
Leu Ile Met Ala Asn Val Ile Leu Leu Pro Val Val Ile Thr Phe Phe	225	230	235
His Asn Lys Asp Met Ser Thr Gly Trp Leu Ile Phe Asn Cys Phe Ser	245	250	255
Asp Thr Phe Phe Ile Leu Asp Leu Ile Cys Asn Phe Arg Thr Gly Ile	260	265	270
Met Asn Pro Lys Ser Ala Glu Gln Val Ile Leu Asn Pro Arg Gln Ile	275	280	285
Ala Tyr His Tyr Leu Arg Ser Trp Phe Ile Ile Asp Leu Val Ser Ser	290	295	300
Ile Pro Met Asp Tyr Ile Phe Leu Leu Ala Gly Gly Gln Asn Arg His	305	310	315
Phe Leu Glu Val Ser Arg Ala Leu Lys Ile Leu Arg Phe Ala Lys Leu	325	330	335
Leu Ser Leu Leu Arg Leu Leu Arg Leu Ser Arg Leu Met Arg Phe Val	340	345	350
Ser Gln Trp Glu Gln Ala Phe Asn Val Ala Asn Ala Val Ile Arg Ile	355	360	365
Cys Asn Leu Val Cys Met Met Leu Leu Ile Gly His Trp Asn Gly Cys	370	375	380
Leu Gln Tyr Leu Val Pro Met Leu Gln Glu Tyr Pro Asp Gln Ser Trp			

002780" 28504260

385				390				395				400				
Val	Ala	Ile	Asn	Gly	Leu	Glu	His	Ala	His	Trp	Trp	Glu	Gln	Tyr	Thr	
				405					410					415		
Trp	Ala	Leu	Phe	Lys	Ala	Leu	Ser	His	Met	Leu	Cys	Ile	Gly	Tyr	Gly	
				420					425					430		
Lys	Phe	Pro	Pro	Gln	Ser	Ile	Thr	Asp	Val	Trp	Leu	Thr	Ile	Val	Ser	
				435					440					445		
Met	Val	Ser	Gly	Ala	Thr	Cys	Phe	Ala	Leu	Phe	Ile	Gly	His	Ala	Thr	
				450					455					460		
Asn	Leu	Ile	Gln	Ser	Met	Asp	Ser	Ser	Ser	Arg	Gln	Tyr	Arg	Glu	Lys	
				465					470					475		
Leu	Lys	Gln	Val	Glu	Glu	Tyr	Met	Gln	Tyr	Arg	Lys	Leu	Pro	Ser	His	
				485					490					495		
Leu	Arg	Asn	Lys	Ile	Leu	Asp	Tyr	Tyr	Glu	Tyr	Arg	Tyr	Arg	Gly	Lys	
				500					505					510		
Met	Phe	Asp	Glu	Arg	His	Ile	Phe	Arg	Glu	Val	Ser	Glu	Ser	Ile	Arg	
				515					520					525		
Gln	Asp	Val	Ala	Asn	Tyr	Asn	Cys	Arg	Asp	Leu	Val	Ala	Ser	Val	Pro	
				530					535					540		
Phe	Phe	Val	Gly	Ala	Asp	Ser	Asn	Phe	Val	Thr	Arg	Val	Val	Thr	Leu	
				545					550					555		
Leu	Glu	Phe	Glu	Val	Phe	Gln	Pro	Ala	Asp	Tyr	Val	Ile	Gln	Glu	Gly	
				565					570					575		
Thr	Phe	Gly	Asp	Arg	Met	Phe	Phe	Ile	Gln	Gln	Gly	Ile	Val	Asp	Ile	
				580					585					590		
Ile	Met	Ser	Asp	Gly	Val	Ile	Ala	Thr	Ser	Leu	Ser	Asp	Gly	Ser	Tyr	
				595					600					605		
Phe	Gly	Glu	Ile	Cys	Leu	Leu	Thr	Arg	Glu	Arg	Arg	Val	Ala	Ser	Val	
				610					615					620		
Lys	Cys	Glu	Thr	Tyr	Cys	Thr	Leu	Phe	Ser	Leu	Ser	Val	Gln	His	Phe	
				625					630					635		
Asn	Gln	Val	Leu	Asp	Glu	Phe	Pro	Ala	Met	Arg	Lys	Thr	Met	Glu	Glu	
				645					650					655		
Ile	Ala	Val	Arg	Arg	Leu	Thr	Arg	Ile	Gly	Lys	Glu	Ser	Ser	Lys	Leu	
				660					665					670		
Lys	Ser	Arg	Leu	Glu	Ser	Pro	Thr	Ile	Arg	Asp	Thr	Ala	Pro	Leu	Phe	
				675					680					685		
Pro	Ile	Pro	Pro	Asp	Thr	Pro	Ser	Phe	Val	Thr	Asp	Ile	Glu	Lys	Asn	
				690					695					700		
Arg	Phe	Phe	Gly	Asp	Asp	Thr	Asp	Asp	Val	His	Ile	Arg	Thr	Arg	Val	

705

710

715

720

Asp Val Glu Arg Gly Ser His Glu Asn Val Ile Ala Ile Met Asp Gly
 725 730 735

Ser Leu Ser Asp Leu Arg Met Glu Asn Glu Ile Gln Ala Arg Lys Ser
 740 745 750

Ser Ser Gly Lys Arg Arg Lys Phe Gln Gln Gln Thr Thr Glu Leu
 755 760 765

0040500.00400